

# Mark Scheme (Results)

March 2012

GCSE Mathematics (2MB01) Higher  
5MB2H (Non-Calculator) Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our qualifications website at [www.edexcel.com](http://www.edexcel.com). For information about our BTEC qualifications, please call 0844 576 0026, or visit our website at [www.btec.co.uk](http://www.btec.co.uk).

If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Alternatively, you can speak directly to the subject team at Pearson about Edexcel qualifications. Their contact details can be found on this link:

[www.edexcel.com/teachingservices](http://www.edexcel.com/teachingservices)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

March 2012

Publications Code G031140

All the material in this publication is copyright

© Pearson Education Ltd 2012

## NOTES ON MARKING PRINCIPLES

- 1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- 2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- 3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- 4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- 5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- 6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
  - i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*  
Comprehension and meaning is clear by using correct notation and labeling conventions.
  - ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*  
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.
  - iii) *organise information clearly and coherently, using specialist vocabulary when appropriate.*  
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

**7 With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

If there is no answer on the answer line then check the working for an obvious answer.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

**8 Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

**9 Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

**10 Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

**11 Linear equations**

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

**12 Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

**13 Range of answers**

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

**Guidance on the use of codes within this mark scheme**

M1 – method mark

A1 – accuracy mark

B1 – Working mark

C1 – communication mark

QWC – quality of written communication

oe – or equivalent

cao – correct answer only

ft – follow through

sc – special case

dep – dependent (on a previous mark or conclusion)

indep – independent

isw – ignore subsequent working

5MB2H_01				
Question	Working	Answer	Mark	Notes
1	$30 \div 12 = 2.5$  $220 \times 2.5$ $40 \times 2.5$ $150 \times 2.5$ $2 \times 2.5$	550 100 375 5	3	<p>M2 for any one of <math>220 + 220 + 110</math> or <math>40 + 40 + 20</math> or <math>150 + 150 + 75</math> or <math>2 + 2 + 1</math> or 550 or 100 or 375 or 5            A1 for all 4 correct values</p> <p>OR</p> <p>M1 for <math>30 \div 12</math> oe or sight of 2.5            M1(dep) for <math>220 \times '2.5'</math> or <math>40 \times '2.5'</math> or <math>150 \times '2.5'</math> or <math>2 \times '2.5'</math> oe or 550 or 100 or 375 or 5            A1 for all 4 correct values</p> <p>OR</p> <p>M1 for <math>220 \div 12</math> or <math>40 \div 12</math> or <math>150 \div 12</math> or <math>2 \div 12</math> or <math>18\frac{1}{3}</math> or <math>3\frac{1}{3}</math> or 12.5 or <math>\frac{1}{6}</math> seen            M1(dep) for <math>'18\frac{1}{3}' \times 30</math> or <math>'3\frac{1}{3}' \times 30</math> or <math>'12.5' \times 30</math> or <math>'\frac{1}{6}' \times 30</math> or 550 or 100 or 375 or 5            A1 for all 4 correct values</p> <p><b>Continued on next page</b></p>

5MB2H_01					
Question		Working	Answer	Mark	Notes
					<p>OR</p> <p>M1 for M1 for <math>220 \div 2</math> or <math>40 \div 2</math> or <math>150 \div 2</math> or <math>2 \div 2</math> or 110 or 20 or 75 or 1 seen  M1(dep) for '110' <math>\times 5</math> or '20' <math>\times 5</math> or '75' <math>\times 5</math> or '1' <math>\times 5</math> or 550 or 100 or 375 or 5 seen  A1 for all 4 correct values</p> <p>(SC: B2 for 275, 50, 187.5 and 2.5  or B1 for 275 or 50 or 187.5 or 2.5, if M0 scored)</p>
2	(a)		Terms must end in 2 or 7	1	B1 for any correct reason; eg terms must end in 2 or 7
	(b)		$5n - 3$	2	B2 for $5n - 3$ (oe, including unsimplified) (B1 for $5n + k$ , $k \neq -3$ or $k$ is absent, or $n = 5n - 3$ )
3		$\left(\frac{1+4}{2}, \frac{2+0}{2}\right)$	(2.5, 1)	2	<p>M1 for <math>\frac{1+4}{2}</math> and <math>\frac{2+0}{2}</math> or for either the <math>x</math> coordinate correct or the <math>y</math> coordinate correct  A1 for (2.5, 1) oe</p> <p>SC: B1 for an answer of (1, 2.5) if M0 scored</p>


5MB2H_01				
Question	Working	Answer	Mark	Notes
4*	$500 \times 1000 \times 400$ $= 200\,000\,000$ $20 \times 50 \times 40 = 40\,000$ $200\,000\,000 \div 40\,000 = 5000$  OR  $(500 \div 20) \times (1000 \div 50) \times$ $(400 \div 40)$ $= 25 \times 20 \times 10 = 5000$	Proof	4	<p>B1 for a correct unit conversion, could be seen on the diagram or in working  M1 for <math>500 \times 1000 \times 400</math> or <math>200\,000\,000</math>  or <math>20 \times 50 \times 40</math> or <math>40\,000</math>  or <math>5 \times 10 \times 4</math> or <math>200</math>  or <math>0.2 \times 0.5 \times 0.4</math> or <math>0.04</math>  M1(dep) for '200 000 000' <math>\div</math> '40 000'  C1 for fully correct working leading to final answer of 5000</p> <p>OR</p> <p>B1 for a correct unit conversion, could be seen on the diagram or in working  M1 for <math>(500 \div 20)</math> or <math>(1000 \div 50)</math> or <math>(400 \div 40)</math> or at least two of 25, 20, 10 seen  M1(dep) for '25' <math>\times</math> '20' <math>\times</math> '10'  C1 for fully correct working leading to final answer of 5000</p>



5MB2H_01				
Question	Working	Answer	Mark	Notes
5	(a)		2	B2 for $8e - 5f$ oe (B1 for $8e$ or $-5f$ )
	(b)		1	B1 cao
	(c)	$3 + 2p - 2$	2	M1 for $2 \times p$ and $2 \times -1$ oe within at most 3 terms seen A1 cao
	(d)	$x(a + b) + y(a + b)$	2	M1 for $x(a + b)$ or $y(a + b)$ or $a(x + y)$ or $b(x + y)$ seen A1 for $(a + b)(x + y)$ oe 2-bracketed expression

5MB2H_01				
Question	Working	Answer	Mark	Notes
6	$30000 \times 20 \div 100 = 6000$ $30000 - 6000 = 24000$ $24000 \div 12$  OR $30000 \div 12 = 2500$ $2500 \times 20 \div 100 = 500$ $2500 - 500$  OR $100 - 20 = 80$ $30000 \times 80 \div 100 = 24000$ $24000 \div 12$  OR $30000 \div 12 = 2500$ $2500 \times 80 \div 100$	2000	3	M1 for $30000 \times 20 \div 100$ oe (= 6000) M1 for $(30000 - '6000') \div 12$ A1 cao  OR  M1 for $30000 \div 12$ (= 2500) M1 for $'2500' - '2500' \times 20 \div 100$ oe A1 cao  OR  M1 for $100 - 20$ (= 80) M1 for $30000 \times 80 \div 100$ oe and dividing by 12 A1 cao  OR  M1 for $30000 \div 12$ (= 2500) M1 for $'2500' \times 80 \div 100$ oe A1 cao

5MB2H_01																														
Question	Working						Answer	Mark	Notes																					
7	$10 \times 8/5 = 16$ $60 - 16$  OR $60 \times 5/8 = 37.5$ $37.5 - 10$						44 km or 27.5 miles	3	B1 for 5 miles = 8 km or equivalent statement or use of conversion factor 8/5 or 5/8 M1 for $10 \times '8/5'$ (= 16) or $60 \times '5/8'$ (= 37.5) A1 for 44 km or 27.5 miles, accept 76 km or 47.5 miles (units must be quoted)																					
8	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>-2</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>y</td> <td>-</td> <td>-6</td> <td>-2</td> <td>2</td> <td>6</td> <td>10</td> </tr> <tr> <td></td> <td>10</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table> OR  Using $y = mx + c$  gradient = 4 y intercept = -2						x	-2	-1	0	1	2	3	y	-	-6	-2	2	6	10		10						Straight line from (-2, -10) to (3, 10)	3	<b>(Table of values)</b> M1 for at least 2 correct attempts to find points by substituting values of x M1 (dep) ft for plotting at least 2 of their points (any points plotted from their table must be correctly plotted) A1 for correct line between $x = -2$ and $x = 3$  <b>(No table of values)</b> M2 for at least 2 correct points and no incorrect points plotted OR line segment of $y = 4x - 2$ drawn (ignore any additional incorrect segments) (M1 for at least 3 correct points with no more than 2 incorrect points) A1 for correct line between $x = -2$ and $x = 3$  <b>(Use of <math>y = mx + c</math>)</b> M2 line segment of $y = 4x - 2$ drawn (ignore any additional incorrect segments) (M1 for line drawn with gradient of 4 OR line drawn with y intercept of -2 and a positive gradient) A1 for correct line between $x = -2$ and $x = 3$
x	-2	-1	0	1	2	3																								
y	-	-6	-2	2	6	10																								
	10																													

5MB2H_01					
Question		Working	Answer	Mark	Notes
9	(a)		6 by 4 rectangle drawn	2	B2 for a 6 by 4 rectangle drawn (B1 for a rectangle drawn with one correct dimension)
	(b)		3-D sketch 	2	M1 for an attempt at a 3-D sketch with a trapezoidal face A1 for a correct 3-D sketch

5MB2H_01				
Question	Working	Answer	Mark	Notes
10*	<p>Angle <math>ABE = 40</math> (vertically opposite angles are equal)            Angle <math>BAE = \text{angle } BEA</math>  <math>= (180 - 40)/2 = 70</math>            (base angles of an isos triangle are equal)  <math>x = 70</math> (alternate angles on parallel lines are equal)</p> <p>OR</p> <p>Angle <math>ABE = 40</math> (vertically opposite angles are equal)            Angle <math>BAE = \text{angle } BEA</math>  <math>= (180 - 40)/2 = 70</math> (base angles of an isosceles triangle are equal)            Angle <math>BEF = 40</math>            (corresponding angles are equal)  <math>x = 180 - 70 - 40 = 70</math>            (angles as a straight line add up to <math>180^\circ</math>)</p>	70	5	<p>B1 for angle <math>ABE = 40</math>, could be marked on the diagram            M1 for <math>(180 - '40')/2 (= 70)</math>            A1 for <math>70^\circ</math> identified as the angle <math>x^\circ</math></p> <p>C2 for fully correct reasons:            'vertically <u>opposite angles</u> are equal' or '<u>vertically opposite angles</u> are equal'            'base <u>angles</u> of an <u>isosceles</u> triangle are <u>equal</u>'            '<u>alternate angles</u> on parallel lines are equal'            (C1 for just one correct reason quoted)</p> <p>OR</p> <p>B1 for angle <math>ABE = 40</math> or angle <math>BEF = 40</math>, could be marked on the diagram            M1 for <math>(180 - '40')/2 (= 70)</math>            A1 for <math>70^\circ</math> identified as the angle <math>x^\circ</math>            C2 for fully correct reasons:            'vertically <u>opposite angles</u> are equal' or '<u>vertically opposite angles</u> are equal'            'base <u>angles</u> of an <u>isosceles</u> triangle are <u>equal</u>'            '<u>corresponding angles</u> on parallel lines are equal'            '<u>angles</u> on a <u>straight line</u> add up to <u>180</u>'            (C1 for just one correct reason quoted)</p>

5MB2H_01				
Question	Working	Answer	Mark	Notes
11	(a)	1	1	B1 cao
	(b)	3	1	B1 cao
	(c)	$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ oe
12	<p>Angle <math>PTO = \text{angle } PSO = 90</math>            Angle <math>TPS = 24 \times 2 = 48</math>  <math>360 - 90 - 90 - 48</math></p> <p>OR</p> <p>Angle <math>PTO = 90</math>            Angle <math>TOP = 180 - 90 - 24</math>  <math>= 66</math>  <math>66 \times 2</math></p>	132	3	<p>M1 for angle <math>PTO = 90</math> or angle <math>PSO = 90</math>, could be marked on the diagram            M1 for <math>360 - 90 - 90 - (24 \times 2)</math>            A1 cao</p> <p>OR</p> <p>M1 for angle <math>PTO = 90</math> or angle <math>PSO = 90</math>, could be marked on the diagram            M1 for <math>2 \times (180 - 90 - 24)</math>            A1 cao</p>

5MB2H_01				
Question	Working	Answer	Mark	Notes
13*	<p>Rectangle – unshaded triangle</p> $(x + 6)(3x - 5) - \frac{1}{2} \times 2x(3x - 5)$ $= 3x^2 + 18x - 5x - 30 - (3x^2 - 5x)$ $= 3x^2 + 18x - 5x - 30 - 3x^2 + 5x \text{ QED}$ <p>OR</p> $(x + 6)(3x - 5) - \frac{1}{2} \times 2x(3x - 5)$ $= (x + 6)(3x - 5) - x(3x - 5)$ $= (3x - 5)(x + 6 - x)$ $= 6(3x - 5) = 18x - 30 \text{ QED}$ <p>OR</p> <p>Shaded trapezium + shaded triangle</p> $\frac{1}{2}(x + 6 - 2x + x + 6)(3x - 5)$ $= 6(3x - 5) = 18x - 30 \text{ QED}$	Proof	4	<p>M1 for using two lengths to find an area  M1(dep) for <math>'(x + 6)(3x - 5)' - \frac{1}{2} \times 2x(3x - 5)'</math>  M1 for <math>3x^2 + 18x - 5x - 30</math> or <math>\frac{1}{2} \times (6x^2 - 10x)</math> or <math>3x^2 - 5x</math>  C1 for a correct completion of the proof resulting in <math>18x - 30</math> from fully correct working</p> <p>OR</p> <p>M1 for using two lengths to find an area  M1(dep) for <math>'(x + 6)(3x - 5)' - \frac{1}{2} \times 2x(3x - 5)'</math>  M1 for factorising process with <math>(3x - 5)</math> as the common factor  C1 for a correct completion of the proof resulting in <math>18x - 30</math> from fully correct working</p> <p>OR</p> <p>M1 for <math>x + 6 - 2x (= 6 - x)</math>  M2 for <math>\frac{1}{2}(x + 6 - 2x + x + 6)(3x - 5)</math>  C1 for a correct completion of the proof resulting in <math>18x - 30</math> from fully correct working</p>

5MB2H_01				
Question	Working	Answer	Mark	Notes
14	$\begin{array}{r} \cdot \\ 0.25 = 0.2555\dots \\ 0.2555\dots \times 10 = 2.5555\dots \\ 0.2555\dots \times 100 = 25.5555\dots \\ 25.5555\dots - 2.5555\dots = 23 \end{array}$ <p><b>Alternative Method</b></p> $\begin{array}{l} x = 0.25555\dots \\ 10x = 2.5555\dots \\ 100x = 25.5555\dots \\ 90x = 23 \end{array}$	$\frac{23}{90}$	3	<p>M1 for <math>0.25555\dots</math> or <math>0.2 + 0.05555\dots</math>  M1 for two correct recurring decimals that, when subtracted, would result a terminating decimal, and attempting the subtraction or <math>\frac{2.3}{9}</math> or <math>\frac{253}{990}</math></p> <p>A1 cao</p> <p><b>Alternative Method</b></p> <p>M1 for <math>x = 0.25555\dots</math>  M1 for two correct recurring decimals that, when subtracted, would result a terminating decimal, e.g. <math>10x = 2.5555\dots</math>, <math>100x = 25.5555\dots</math> and attempting the subtraction or <math>9x = 2.3</math> or <math>90x = 23</math> or <math>900x = 230</math> or <math>\frac{2.3}{9}</math> or <math>\frac{253}{990}</math></p> <p>A1 cao</p> <p>SC: B1 for fraction with denominator 90 if M0</p>
15	$\begin{array}{l} \text{Grad} = \frac{1}{3} \\ 3 = 6 \times \frac{1}{3} + c \\ c = 1 \end{array}$	$y = \frac{1}{3}x + 1$	3	<p>M1 for <math>-3 \times m = -1</math> oe or <math>\frac{1}{3}</math> seen</p> <p>M1 for <math>3 = 6 \times \frac{1}{3} + c</math> or <math>y - 3 = \frac{1}{3}(x - 6)</math></p> <p>A1 for <math>y = \frac{1}{3}x + 1</math> oe</p>



5MB2H_01				
Question	Working	Answer	Mark	Notes
16	(a) $\frac{15}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$	$3\sqrt{5}$	2	M1 for $\frac{15}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$ A1 for $\frac{15\sqrt{5}}{5}$ or better
	(b) $(1 + \sqrt{3})(1 + \sqrt{3})$ $= 1 + \sqrt{3} + \sqrt{3} + 3$ $= 4 + 2\sqrt{3}$	4, 2	2	M1 for $1 \times 1 + 1 \times \sqrt{3} + 1 \times \sqrt{3} + \sqrt{3} \times \sqrt{3}$ oe A1 cao
17	$\frac{3(a-b)+2b}{b(a-b)}$ $\frac{3a-3b+2b}{b(a-b)}$	$\frac{3a-b}{b(a-b)}$	3	M1 for a common denominator of $b(a-b)$ oe M1 for $\frac{3(a-b)+2b}{'b(a-b)'} \text{ or } \frac{3(a-b)}{'b(a-b)'} + \frac{2b}{'b(a-b)'}$ A1 for $\frac{3a-b}{b(a-b)}$ or $\frac{3a-b}{ba-b^2}$



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UG031140 March 2012

For more information on Edexcel qualifications, please visit  
[www.edexcel.com/quals](http://www.edexcel.com/quals)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

*Downloaded from TopLevels.co.uk*

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

