

3.




Figure 1

Figure 1 shows a sketch of part of the curve with equation $y = 4x - xe^{\frac{1}{2}x}$, $x \geq 0$

The curve meets the x -axis at the origin O and cuts the x -axis at the point A .

(a) Find, in terms of $\ln 2$, the x coordinate of the point A . (2)

(b) Find

$$\int xe^{\frac{1}{2}x} dx$$
(3)

The finite region R , shown shaded in Figure 1, is bounded by the x -axis and the curve with equation

$$y = 4x - xe^{\frac{1}{2}x}, \quad x \geq 0$$

(c) Find, by integration, the exact value for the area of R .
Give your answer in terms of $\ln 2$ (3)



8.




Diagram not to scale

Figure 3

Figure 3 shows a sketch of part of the curve C with equation

$$y = 3^x$$

The point P lies on C and has coordinates $(2, 9)$.

The line l is a tangent to C at P . The line l cuts the x -axis at the point Q .

(a) Find the exact value of the x coordinate of Q .

(4)

The finite region R , shown shaded in Figure 3, is bounded by the curve C , the x -axis, the y -axis and the line l . This region R is rotated through 360° about the x -axis.

(b) Use integration to find the exact value of the volume of the solid generated.

Give your answer in the form $\frac{p}{q}$ where p and q are exact constants.

[You may assume the formula $V = \frac{1}{3}\pi r^2 h$ for the volume of a cone.]

(6)



