

Mark Scheme

Pearson Edexcel GCSE (9-1)
Mathematics – 1MA1
Trial of Specimen Papers (Set 1)

Paper 2 (1MA1/2H): Calculator
Higher Tier

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

March 2016

All the material in this publication is copyright

© Pearson Education Ltd 2016

General marking guidance

These notes offer general guidance, but the specific notes for examiners appertaining to individual questions take precedence.

- 1** All candidates must receive the same treatment. Examiners must mark the last candidate in exactly the same way as they mark the first.

Where some judgement is required, mark schemes will provide the principles by which marks will be awarded; exemplification/indicative content will not be exhaustive. When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the response should be sent to review.

- 2** All the marks on the mark scheme are designed to be awarded; mark schemes should be applied positively. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme. If there is a wrong answer (or no answer) indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

Questions where working is not required: In general, the correct answer should be given full marks.

Questions that specifically require working: In general, candidates who do not show working on this type of question will get no marks – full details will be given in the mark scheme for each individual question.

- 3** **Crossed out work**

This should be marked **unless** the candidate has replaced it with an alternative response.

- 4** **Choice of method**

If there is a choice of methods shown, mark the method that leads to the answer given on the answer line.

If no answer appears on the answer line, mark both methods **then award the lower number of marks.**

- 5** **Incorrect method**

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review for your Team Leader to check.

6 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working as you can check the answer, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question or its context. (eg. an incorrectly cancelled fraction when the unsimplified fraction would gain full marks).

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect (eg. incorrect algebraic simplification).

8 Probability

Probability answers must be given as a fraction, percentage or decimal. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

9 Linear equations

Unless indicated otherwise in the mark scheme, full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously identified in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

10 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and all numbers within the range.

Guidance on the use of abbreviations within this mark scheme

- M** method mark awarded for a correct method or partial method
- P** process mark awarded for a correct process as part of a problem solving question
- A** accuracy mark (awarded after a correct method or process; if no method or process is seen then full marks for the question are implied but see individual mark schemes for more details)
- C** communication mark
- B** unconditional accuracy mark (no method needed)
- oe** or equivalent
- cao** correct answer only
- ft** follow through (when appropriate as per mark scheme)
- sc** special case
- dep** dependent (on a previous mark)
- indep** independent
- awrt** answer which rounds to
- isw** ignore subsequent working

Mark scheme GCSE (9 – 1) Mathematics

Paper 1MA1_2H			
Question	Working	Answer	Notes
1		$t = \frac{w - 11}{3}$	<p>M1 For isolating term in t, eg. $3t = w - 11$ or dividing all terms by 3, eg. $\frac{w}{3} = \frac{3t}{3} + \frac{11}{3}$</p> <p>A1 for $t = \frac{w-11}{3}$ oe</p>
2	<p>£: $1980 \div 1.34 = 1477.61$ $2250 \div 1.52 =$ 1480.26</p> <p>€: $1480 \times 1.34 = 1983.2$ $2250 \div 1.52 \times 1.34 = 1983.55$</p> <p>\$: $1480 \times 1.52 = 2249.6$ $1980 \div 1.34 \times 1.52 =$ 2245.9</p>	Jardins of Paris	<p>P1 correct process to convert one price to another currency, eg $1980 \div 1.34$</p> <p>P1 for a complete process leading to 3 prices in the same currency</p> <p>C1 for 3 correct and consistent results and a correct comparison made.</p>
3		Mean of 96 or net deviation of 0 so target met	<p>M1 for correct interpretation of the graph, with at least one correct reading or a line drawn through 96 with at least one correct deviation</p> <p>M1 complete method to find mean of six months sales, eg. $(110+84+78+94+90+120) \div 6 (= 96)$ or the sum of six deviations, eg. $(14-12-18-2-6+24) \div 6 (= 0)$</p> <p>C1 for a correct answer of 96 or 0 with correct conclusion</p>

Paper 1MA1_2H				
Question	Working	Answer	Notes	
4	(a)	$160 < h \leq 170$	B1	for identifying the correct class interval
	(b)	1. Points should be plotted at mid-interval values 2. The polygon should not be closed	C1 C1	for a correct error identified for a correct error identified
5	(a)	graph	M1 C1 C1	for method to start to find distance cycled in 36 mins, eg. line drawn of correct gradient or $15 \times \frac{36}{60}$ or 15×36 for correct graph from 9.00 am to 9.36 am for graph drawn from "(9.36, 9)" to (10.45, "9" + 8)
	(b)	4.5	M1 A1	for 18×0.25 oe cao
6		8112	M1 A1	for complete method, eg. 7500×1.04^2 cao

Paper 1MA1_2H			
Question	Working	Answer	Notes
7		No with supporting evidence	<p>P1 for the start of a correct process, eg. two of x, $2x$ and $2x+7$ or a fully correct trial, eg. $5 + 10 + 17 = 32$</p> <p>P1 (dep on P1) for setting up an equation using 3 algebraic terms, eg. $x + 2x + 2x + 7 = 57$ or a correct trial totalling 57, eg. $10 + 20 + 27 = 57$</p> <p>C1 for a correct deduction from correct answers, eg. Chris has 20 so it is impossible for all to have 20 since 60 marbles would be needed.</p>
8		66.9	<p>P1 for process to find the area of one shape, eg. $19 \times 16 (= 304)$ or $\pi \times 8^2 (= 201.06\dots)$</p> <p>P1 for process to find the shaded area, eg. $"304" - "201.06" \div 2 (= 203.46\dots)$</p> <p>P1 for a complete process to find required percentage, eg. $\frac{"203.46"}{304} \times 100$</p> <p>A1 for answer in range 66 to 68</p>
9		135	<p>B1 for identifying the angle of 70° (on the diagram), showing understanding of notation</p> <p>P1 for process to find an angle in triangle ABC, eg. for process to find angle BAC, eg. $(180 - 50) \div 2 (= 65^\circ)$</p> <p>A1 for 135</p>

Paper 1MA1_2H			
Question	Working	Answer	Notes
10 (a)		-1.5	M1 for method to find gradient, eg. $210 \div 140$ A1 for correct interpretation of the negative gradient
(b)			C1 for explanation, eg. rate of change of depth of water in tank
11 (a)		0.49	M1 for 0.7×0.7 A1 for 0.49 oe
(b)		0.51	M1 for a correct process, eg. $1 - "0.49"$ or $0.7 \times 0.3 + 0.3 \times 0.7 + 0.3 \times 0.3$ A1 for 0.51 oe
12 (a)		0.4	B1 For 0.4 oe
(b)		0.586	M1 for "3.48207....." \div 17.34 or $3.48207..... \div "17.34"$ or 0.200811... A1 for 0.585 to 0.586
13	$(3x - 1)(4x^2 + 20x - 3x - 15)$ $(x + 5)(12x^2 - 4x - 9x + 3)$ $(4x - 3)(3x^2 - x + 15x - 5)$	Fully correct algebra to show given result	M1 for method to find the product of any two linear expressions; eg. 3 correct terms or 4 terms ignoring signs M1 (dep) for method of 6 products, 4 of which are correct (ft their first product) A1 for fully accurate working to give the required result

Paper 1MA1_2H			
Question	Working	Answer	Notes
14	angle $BAD =$ angle $DCA = 22.62^\circ$ angle $DBA =$ angle $DAC = 67.38^\circ$	33.8	P1 for recognition of similar triangles or equal ratio of sides OR for a method to find angle BAD or angle DBA and state that this is the same as angle DCA or angle DAC P1 for process to find CB , eg. $\frac{5}{13} = \frac{13}{CB}$ A1 for an answer rounding to 33.8
15		18.3	P1 for a start to the process interpreting the information correctly, eg. $T = k\sqrt{L}$ oe P1 for a correct scale factor of $\sqrt{1.4}$ A1 for 18.3 to 18.4
16		84	M1 for correct interpretation of given information leading to a method to find fd, eg. $20 \div 100$ (thousand) or for an acceptable key P1 for a process to find at least two required frequencies, eg. $0.8 \times 50 (= 40)$, $0.6 \times 50 (= 30)$, $0.14 \times 100 (= 14)$ A1 for 84 cao

Paper 1MA1_2H			
Question	Working	Answer	Notes
17		$n^2 - n + 1$ oe	<p>M1 for correct deduction from differences, eg. 2nd difference of 2 implies $1n^2$ or sight of $1^2, 2^2, 3^2, \dots$</p> <p>M1 for sight of $1^2, 2^2, 3^2, \dots$ linked with 1, 2, 3, ...</p> <p>A1 for $n^2 - n + 1$ oe</p> <p>OR</p> <p>M1 for $a + b + c = 1$ or $4a + 2b + c = 3$ or $9a + 3b + c = 7$ oe</p> <p>M1 for a method to eliminate one unknown leaving simultaneous equations in the other two</p> <p>A1 for $n^2 - n + 1$ oe</p>
18		$3x^2 + 10x$	<p>M1 start a chain of reasoning, eg. $3(x+2)^2 - 2(x+2) - 8$</p> <p>M1 continue chain by expanding brackets correctly, eg. $3x^2 + 12x + 12 - 2x - 4 - 8$</p> <p>A1 for $3x^2 + 10x$ ($a = 3, b = 10$)</p>
19		8.63 to 8.65	<p>P1 for a start of process, eg. $0.5x(x - 2) = 2.5$</p> <p>P1 for rearranging to give a quadratic equation, eg. $x^2 - 2x - 5 (= 0)$ oe.</p> <p>P1 (dep on P1) for a process to solve their 3-term quadratic equation, condoning one sign error in use of formula ($x = 3.449\dots$ and $x = -1.449\dots$)</p> <p>P1 for selecting the positive value of x and applying Pythagoras to find the hypotenuse, eg. $\sqrt{("3.449")^2 + ("1.449")^2}$ ($= 3.74\dots$)</p> <p>P1 (dep on previous P1) for complete process to find perimeter</p> <p>A1 for answer in the range 8.63 to 8.65</p>

Paper 1MA1_2H			
Question	Working	Answer	Notes
20 (a)		3 to 4	C1 for a tangent drawn at $t = 6$ B1 for a gradient in the range 3 to 4 or ft "tangent"
(b)		452	C1 for splitting the area into 3 strips and a method of finding the area of one shape under the graph, eg. $\frac{1}{2} \times 4 \times 35 (= 70)$ M1 for complete process to find the area under the graph, eg " $70 + \frac{1}{2} \times 4 \times (35 + 51) (= 172) + \frac{1}{2} \times 4 \times (51 + 54) (= 210) [= 452]$ " A1 for 452
21		10169 or 10171	P1 for correct use of formula to find number in 2016, eg. $1.05(9500 - 250) (= 9712.5)$ for complete iterative process, P1 eg. 2017: $1.05("9712.5" - 250) (= 9935.625)$ 2018: $1.05("9935.625" - 250)$ for answer of 10169.90... rounded or truncated to nearest whole number C1
22		1.5	B1 for any correct bound clearly identified, eg. $99.65 \rightarrow x \rightarrow 99.75$ or $66.5 \rightarrow y \rightarrow 67.5$ M1 (dep on B1) for method to find UB, eg. " $99.75 \div 66.5$ " A1 for 1.5
23		$y = -\frac{4}{3}x + \frac{25}{3}$ oe	M1 for method to find gradient of tangent, eg. $-1 \div \frac{3}{4} = -\frac{4}{3}$

Paper 1MA1_2H			
Question	Working	Answer	Notes
			<p>M1 (dep) for method to find y-intercept using $y = -\frac{4}{3}x + c$</p> <p>A1 $y = -\frac{4}{3}x + \frac{25}{3}$ oe</p>
24		Proof	<p>C1 for joining AO (extended to D) and considering angles in two triangles (algebraic notation may be used here)</p> <p>C1 for using isosceles triangle properties to find angle BOD (eg. $x + x = 2x$) or angle COD (eg. $y + y = 2y$)</p> <p>C1 for angle $BOC = 2x + 2y$ [= $2 \times \text{angle } BAO + 2 \times \text{angle } CAO$]</p> <p>C1 for completion of proof with all relevant reasons given, eg. base <u>angles</u> of <u>isosceles</u> triangle are <u>equal</u> and sum of <u>angles</u> at a <u>point</u> is <u>360°</u></p>

