

2.




Figure 1

The uniform lamina $OABCD$, shown in Figure 1, is formed by removing the triangle OAD from the square $ABCD$ with centre O . The square has sides of length $2a$.

(a) Show that the centre of mass of $OABCD$ is $\frac{2}{9}a$ from O . (4)

The mass of the lamina is M . A particle of mass kM is attached to the lamina at D to form the system S . The system S is freely suspended from A and hangs in equilibrium with AO vertical.

(b) Find the value of k . (4)



4. A ladder AB , of weight W and length $2l$, has one end A resting on rough horizontal ground. The other end B rests against a rough vertical wall. The coefficient of friction between the ladder and the wall is $\frac{1}{3}$. The coefficient of friction between the ladder and the ground is μ . Friction is limiting at both A and B . The ladder is at an angle θ to the ground, where $\tan \theta = \frac{5}{3}$. The ladder is modelled as a uniform rod which lies in a vertical plane perpendicular to the wall.

Find the value of μ .

(9)



Question 4 continued

Lined area for writing the answer to Question 4.



Question 4 continued

Lined area for writing the answer to Question 4.

(Total 9 marks)

Q4



5.




Figure 2

A particle P of mass 10 kg is projected from a point A up a line of greatest slope AB of a fixed rough plane. The plane is inclined at angle α to the horizontal, where $\tan \alpha = \frac{5}{12}$ and $AB = 6.5$ m, as shown in Figure 2. The coefficient of friction between P and the plane is μ . The work done against friction as P moves from A to B is 245 J.

(a) Find the value of μ . **(5)**

The particle is projected from A with speed 11.5 m s^{-1} . By using the work-energy principle,

(b) find the speed of the particle as it passes through B . **(4)**



Question 5 continued

Lined writing area for the question response.

(Total 9 marks)

Q5

Grading box for Q5



7.




Figure 3

At time $t = 0$, a particle is projected from a fixed point O on horizontal ground with speed $u \text{ m s}^{-1}$ at an angle θ° to the horizontal. The particle moves freely under gravity and passes through the point A when $t = 4 \text{ s}$. As it passes through A , the particle is moving upwards at 20° to the horizontal with speed 15 m s^{-1} , as shown in Figure 3.

- (a) Find the value of u and the value of θ . **(7)**

At the point B on its path the particle is moving downwards at 20° to the horizontal with speed 15 m s^{-1} .

- (b) Find the time taken for the particle to move from A to B . **(2)**

The particle reaches the ground at the point C .

- (c) Find the distance OC . **(3)**



Question 7 continued

Lined area for writing the answer to Question 7 continued.



8. Three identical particles P , Q and R , each of mass m , lie in a straight line on a smooth horizontal plane with Q between P and R . Particles P and Q are projected directly towards each other with speeds $4u$ and $2u$ respectively, and at the same time particle R is projected along the line away from Q with speed $3u$. The coefficient of restitution between each pair of particles is e . After the collision between P and Q there is a collision between Q and R .

(a) Show that $e > \frac{2}{3}$ **(7)**

It is given that $e = \frac{3}{4}$

(b) Show that there will not be a further collision between P and Q . **(6)**



