# 

# GCSE Mathematics

Paper 2 Higher Tier

Mark scheme

8300 November 2017

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Copyright © 2017 AQA and its licensors. All rights reserved.

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

#### **Glossary for Mark Schemes**

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

| М               | Method marks are awarded for a correct method which could lead to a correct answer.  |
|-----------------|--|
| Α               | Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied. |
| В               | Marks awarded independent of method.   |
| ft              | Follow through marks. Marks awarded for correct working following a mistake in an earlier step.  |
| SC              | Special case. Marks awarded for a common misinterpretation which has some mathematical worth.  |
| M dep           | A method mark dependent on a previous method mark being awarded.   |
| B dep           | A mark that can only be awarded if a previous independent mark has been awarded.   |
| oe              | Or equivalent. Accept answers that are equivalent.   |
|                 | eg accept 0.5 as well as $\frac{1}{2}$   |
| [a, b]          | Accept values between a and b inclusive.   |
| [a, b)          | Accept values a ≤ value < b  |
| 3.14            | Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416   |
| Use of brackets | It is not necessary to see the bracketed work to award the marks.  |

Examiners should consistently apply the following principles

#### Diagrams

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

#### Responses which appear to come from incorrect methods

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

#### Questions which ask students to show working

Instructions on marking will be given but usually marks are not awarded to students who show no working.

#### Questions which do not ask students to show working

As a general principle, a correct response is awarded full marks.

#### Misread or miscopy

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

#### Further work

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

#### Choice

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

#### Work not replaced

Erased or crossed out work that is still legible should be marked.

#### Work replaced

Erased or crossed out work that has been replaced is not awarded marks.

#### Premature approximation

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

#### **Continental notation**

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the student intended it to be a decimal point.

| Question | Answer              | Mark | Comments |  |
|----------|---------------------|------|----------|--|
|          | <u>31</u><br>8      | B1   |          |  |
| 1        | Additional Guidance |      |          |  |
|          |                     |      |          |  |

|   | 250%                  | B1 |  |  |  |  |  |
|---|-----------------------|----|--|--|--|--|--|
| 2 | 2 Additional Guidance |    |  |  |  |  |  |
|   |                       |    |  |  |  |  |  |

| 3 | $\left(\frac{1}{3}, \frac{1}{9}\right)$ B1 |
|---|--|
|   | Additional Guidance                        |
|   |  |

|   | kg/m <sup>3</sup> | B1         |          |  |
|---|-------------------|------------|----------|--|
| 4 | Ade               | ditional G | Buidance |  |
|   |                   |            |          |  |

| Question | Answer  | Mark | Comments   |  |  |
|----------|---|------|--|--|--|
|          | Alternative method 1  |      |  |  |  |
| -        | 12 <i>x</i> – 8   | M1   | May be seen in a grid  |  |  |
|          | their $12x - 2x = -5$ + their 8<br>or $10x = 3$<br>or their $-8 + 5 = 2x$ - their $12x$<br>or $-3 = -10x$ | M1   | Collecting two terms in $x$ and two<br>constant terms correctly<br>oe eg $10x - 3 = 0$ |  |  |
|          | 0.3 or $\frac{3}{10}$   | A1ft | ft M1M0 or M0M1 with exactly one error   |  |  |
|          | Alternative method 2  |      |  |  |  |
| 5        | $\frac{x}{2} - \frac{5}{4}$   | M1   |  |  |  |
|          | $3x - \text{their } \frac{x}{2} = \text{their } -\frac{5}{4} + 2$   |      | Collecting two terms in <i>x</i> and two constant terms correctly                      |  |  |
|          | or $\frac{5}{2}x = \frac{3}{4}$   | M1   | oe eg $\frac{5}{2}x - \frac{3}{4} = 0$   |  |  |
|          | or $-2$ + their $\frac{5}{4}$ = their $\frac{x}{2} - 3x$  |      |  |  |  |
| -        | or $-\frac{3}{4} = -\frac{5}{2}x$   |      |  |  |  |
|          | 0.3 or $\frac{3}{10}$   | A1ft | ft M1M0 or M0M1 with exactly one error   |  |  |

|   | Additional Guidance  |          |
|---|--|----------|
|   | 12x - 2 = 2x - 5   | MO       |
|   | 10x = -3   | M1       |
|   | x = -0.3   | A1ft     |
|   | 12x - 8 = 2x - 5   | M1       |
|   | 10x = -5   | MO       |
|   | $x = \frac{-5}{10}$  | A1ft     |
|   | 12x - 8 = 2x - 5   | M1       |
|   | 14x = 3  | MO       |
| 5 | $x = \frac{3}{14}$   | A1ft     |
|   | 12x - 8 = 2x - 5   | M1       |
|   | 14x = -13  | MO       |
|   | $x = -\frac{13}{14}$ (two errors)                                      | A0ft     |
|   | 12x - 8 = 8x - 20  | M1M0A0   |
|   | Any ft answer must be exact or rounded or truncated to at least 2 dp   |          |
|   | The last two marks can be implied without the collection of terms seen |          |
|   | eg $12x - 6 = 2x - 5$ and answer 0.1                                   | M0M1A1ft |
|   | Collecting terms before the bracket has been expanded                  | Zero     |

| Question | Answer  | Mark | Comments  |    |  |
|----------|---|------|---|----|--|
|          | Correct product using a point on<br>the curve<br>or<br>correct division using a point on the<br>curve | B1   | eg 2 x 12 (= 24) or 3 x 8 (= 24)<br>or 5 x 4.8 (= 24) or 6 x 4 (= 24)<br>or 10 x 2.4 (= 24) or 24 $\div$ 2 = 12<br>or 24 $\div$ 6 = 4 |    |  |
|          | Additional Guidance   |      |   |    |  |
|          | 1 × 24 (= 24)   |      |   | B0 |  |
| 6(a)     | 12 + 12 (= 24)  |      |   | B0 |  |
|          | $3 \times 4 \times 2 = 24$  |      |   | B0 |  |
|          | For multiplication, 24 does not have to be shown  |      |   |    |  |
|          | Ignore any units seen   |      |   |    |  |
|          | Ignore any lines on the graph   |      |   |    |  |
|          | 8 × 3 = 24 and 12 + 12 = 24 (choice)  |      |   | B0 |  |
|          | area 6 and length 4 and volume 24   |      |   | B0 |  |

| Question | Answer   | Mark  | Comments |  |  |
|----------|--|-------|----------|--|--|
|          | Alternative method 1                             |       |          |  |  |
|          | Reading from 5 on the graph to give [4.7, 4.9]   | M1    |          |  |  |
|          | $\frac{1}{2} \times 6 \times h = [4.7, 4.9]$     | M1dep | oe       |  |  |
|          | or [4.7, 4.9] ÷ $(\frac{1}{2} \times 6)$         |       |          |  |  |
|          | [1.56, 1.64]                                     | A1    |          |  |  |
|          | Alternative method 2                             |       |          |  |  |
|          | 24 ÷ 5 or 4.8 or $\frac{1}{2} \times 6 \times h$ | M1    | oe       |  |  |
| 6(b)     | or $\frac{1}{2} \times 6 \times h \times 5$      |       |          |  |  |
|          | $\frac{1}{2} \times 6 \times h = 24 \div 5$      |       | oe       |  |  |
|          | or $24 \div 5 \div (\frac{1}{2} \times 6)$       |       |          |  |  |
|          | or $\frac{1}{2} \times 6 \times h \times 5 = 24$ | M1dep |          |  |  |
|          | or 15 <i>h</i> = 24                              |       |          |  |  |
|          | or $24 \div (\frac{1}{2} \times 6 \times 5)$     |       |          |  |  |
|          | or 24 ÷ 15                                       |       |          |  |  |
|          | 1.6  | A1    |          |  |  |
|          | Additional Guidance                              |       |          |  |  |
|          |  |       |          |  |  |

| Question | Answer   | Mark | Comments |        |
|----------|--|------|----------|--------|
|          | Enlargement  | B1   |          |        |
|          | Scale factor (x) $\frac{1}{3}$   | B1   |          |        |
| -        | Centre (5, 1)  | B1   |          |        |
|          | Additional Guidance  |      |          |        |
| 7        | Enlarge (x) $\frac{1}{3}$ (5, 1)   |      |          | B1B1B1 |
|          | Reduction or makes bigger or unenlargement or increase or negative enlargement   |      |          | 1st B0 |
|          | Any other transformation mentioned or implied such as reflection, rotation or translation loses the mark for enlargement |      |          |        |
|          | eg enlarged and moved up 4 or enlarged and $\begin{pmatrix} -2 \\ 2 \end{pmatrix}$                                       |      |          | 1st B0 |
|          | Do not accept ÷ 3 for scale factor   |      |          | 2nd B0 |

|   | [0, 5] × 20 + [5, 10] × 48<br>+ [10, 15] × 30 + [15, 20] × 22<br>or 1170 | M1             | Must add 4 products     |      |
|---|--|----------------|-------------------------|------|
|   | their 1170 ÷ 120   | M1dep          |                         |      |
| 8 | 9.75 or $\frac{39}{4}$ or $9\frac{3}{4}$                                 | A1             |                         |      |
|   | Ad   | ditional G     | uidance                 |      |
|   | 1170 ÷ 120 or 9.75 with 5 < <i>x</i> ≤ 10 on                             | answer lin     | e                       | M2A0 |
|   | Do not allow M1 for working in the ta working lines                      | ble if a diffe | erent method is used in |      |

| Question | Answer  | Mark         | Commen  | ts                                  |  |
|----------|---|--------------|---|-------------------------------------|--|
|          | $\tan x = \frac{3}{7} \text{ or } \tan^{-1} \frac{3}{7}$<br>or $\sin x = \frac{3(\sin 90)}{\sqrt{3^2 + 7^2}}$<br>or $\sin x = \frac{3(\sin 90)}{\sqrt{58}}$<br>or $\cos x = \frac{7}{\sqrt{3^2 + 7^2}}$<br>or $\cos x = \frac{7}{\sqrt{58}}$<br>or $\cos x = \frac{7}{\sqrt{58}}$<br>or $90 - \tan^{-1} \frac{7}{3}$<br>or $90 - [66.7, 66.81]$<br>or $90 - 67$ | M1           | oe<br>eg cos $x = \frac{7^2 + (\sqrt{7^2 + 3})}{2 \times \sqrt{3^2 + 7}}$<br>Any letter | $\frac{\overline{7}}{7^2} \times 7$ |  |
| 9        | [23, 23.3]  | A1           |   |                                     |  |
|          | Additional Guidance   |              |   |                                     |  |
|          | $\tan = \frac{3}{7}$ or $\tan \frac{3}{7}$ or $\tan^{-1} = \frac{3}{7}$ (u  | MO           |   |                                     |  |
|          | Answer [23, 23.3] (possibly coming f  | rom scale    | drawing)  | M1A1                                |  |
|          | If using sine rule must rearrange to  | $\sin x = f$ | or M1   |                                     |  |
|          | If using cosine rule must rearrange to  |              |   |                                     |  |
|          | Allow [0.42, 0.43] for $\frac{3}{7}$  |              |   |                                     |  |
|          | Allow 2.33 for $\frac{7}{3}$  |              |   |                                     |  |
|          | Allow [7.6, 7.62] for $\sqrt{3^2 + 7^2}$  |              |   |                                     |  |

| Question | Answer  | Mark | Comments |      |  |
|----------|---|------|----------|------|--|
|          | 3 6 9<br>or 23 + 12<br>or $1.5n^2$                                    | M1   |          |      |  |
|          | 35  | A1   |          |      |  |
| 10       | Additional Guidance   |      |          |      |  |
|          | Answer line blank with 35 as next ter                                 | M1A1 |          |      |  |
|          | Answer line has attempt at term to te                                 | M1A0 |          |      |  |
|          | 35 seen on dotted line in sequence but a different answer given eg 50 |      |          | M1A0 |  |

| 11 | $\frac{x^2}{2x^2+1}$ B1 |
|----|-------------------------|
|    | Additional Guidance     |
|    |                         |

| Question | Answer  | Mark | Commen   | ts       |  |
|----------|---|------|--|----------|--|
|          | 64 000 000 ÷ 95 000<br>or 673.() or 674 or $\frac{12\ 800}{19}$<br>or<br>82 000 000 ÷ 140 000<br>or 585.() or 586 or $\frac{4100}{7}$ | M1   | oe population ÷ area<br>Accept a pair of consiste<br>eg 64 ÷ 95 or 0.673 o<br>and 82 ÷ 140 or 0.585  | or 0.674 |  |
|          | 673.() or 674 or 670<br>and<br>585.() or 586 or 590<br>or $\frac{89\ 600}{133}$ and $\frac{77\ 900}{133}$                             | A1   | Correct comparable values from<br>consistent divisions<br>eg 0.674 and 0.586<br>Accept 700 with division seen for UK<br>Accept 600 with division seen for<br>Germany |          |  |
| 12       | Comparable values<br>and correct conclusion   | A1ft | eg<br>673 and 585 and greater for UK<br>0.673 and 0.585 and greater for UK<br>ft M1A0 and comparable values<br>Ignore further work                                   |          |  |
|          | Additional Guidance   |      |  |          |  |
|          | Comparable values means both must be in the same form eg fractions with common denominators   |      |  |          |  |
|          | 64 000 000 ÷ 95 000 = 67.4<br>82 000 000 ÷ 140 000 = 5857<br>Germany is higher  |      | M1<br>A0<br>A1ft   |          |  |
|          | Ignore subtraction of results   |      |  |          |  |
|          | 673 and 585 and UK has more people per square mile  |      |  | M1A1A1ft |  |
|          | 673 and 585 and Germany has more space for their population   |      |  | M1A1A1ft |  |
|          | 673 and 585 and UK's population is less spread out  |      |  | M1A1A1ft |  |
|          | 673 and 585 and UK is more than Germany   |      |  | M1A1A1ft |  |
|          | 673 and 585 and UK is 78 more than Germany (ignore further work)  |      |  | M1A1A1ft |  |

|         | 673 and 585 and the difference is 88   | M1A1A0ft |
|---------|--|----------|
|         | 673 and 585 and UK population is bigger  | M1A1A0ft |
|         | 673 and 586 and UK   | M1A1A0ft |
| 12 cont | 673 and 585 and Germany has more space   | M1A1A0ft |
|         | 673 > 585 (unless links to countries in working)                                       | M1A1A0ft |
|         | $\frac{12\ 800}{19}$ and $\frac{4100}{7}$ and UK is greater (fractions not comparable) | M1A0A0ft |

| Question | Answer   | Mark       | Comme   | ents |  |
|----------|--|------------|---------|------|--|
|          | $\left(-\frac{1}{3},-1\right)$   | B1         |         |      |  |
| 13       | Ado  | ditional G | uidance |      |  |
|          |  |            |         |      |  |
| 14(-)    | $\frac{3}{4} \times \frac{3}{4} \times 15$<br>or<br>$\frac{3}{4} \times 15 \text{ or } 11.25$<br>and $\frac{3}{4} \times \text{their } 11.25$<br>8.4(375) or 8.44 or 8.438<br>or $\frac{135}{16}$ or $8\frac{7}{16}$ | M1<br>A1   | oe      |      |  |
| 14(a)    | Additional Guidance  |            |         |      |  |
| -        | 8.43 or 8.437  |            |         | M1A1 |  |
| _        | 8.4 seen, answer 8   |            |         | M1A1 |  |
|          | $\frac{3}{4}$ of 11.25 (unless correctly evaluated)  |            |         | МО   |  |
|          | $\frac{3}{4}$ × 8.4375, answer 6.328 (further work)  |            |         | M1A0 |  |
|          | 11.25 + 8.4375, answer 19.6875 (furt   | her work)  |         | M1A0 |  |

| Question | Answer  | Mark | Comments  |  |  |
|----------|---|------|---|--|--|
|          | Alternative method 1  |      |   |  |  |
|          | Ticks second box<br>and [7.425, 7.5375]<br>or<br>Ticks second box<br>and correctly evaluates<br>$\frac{2}{3}$ × their 11.25 | B2ft | ft correct box ticked for comparing with<br>their answer to (a)<br>B1ft [7.425, 7.5375]<br>with no or incorrect decision<br>or<br>Correctly evaluates<br>$\frac{2}{3} \times$ their 11.25   |  |  |
|          | Alternative method 2  |      | with no or incorrect decision   |  |  |
| 14(b)    | Ticks second box<br>and valid comparison  | B2   | eg $\frac{8}{12}$ and $\frac{9}{12}$<br>0.66 or 0.67 and 0.75<br>66.()% or 67% and 75%<br>$\frac{9}{16}$ and $\frac{8}{16}$<br>clear diagrams showing $\frac{2}{3}$ and $\frac{3}{4}$<br>B1 Ticks second box<br>and incomplete comparison<br>eg $\frac{8}{12}$ and $\frac{3}{4}$<br>two thirds is less than three quarters<br>$\frac{3}{4} \times \frac{3}{4} = \frac{9}{16}$ and $\frac{3}{4} \times \frac{2}{3} = \frac{6}{12}$<br>or<br>Valid comparison (that would score |  |  |

| Additional Guidance  |      |
|--|------|
| In Alt 1 only follow through their answer to (a) for the comparison, the working for $\frac{2}{3}$ × their 11.25 must be correct |      |
| (a) answer 6.5 (b) Ticks first box and 7.5 seen  | B2ft |
| Accept 0.66 or 0.67 for $\frac{2}{3}$  |      |
| Using 0.6 for $\frac{2}{3}$  | B0   |

| Question | Answer  | Mark  | Comments   |  |  |
|----------|---|-------|--|--|--|
|          | Alternative method 1  |       |  |  |  |
|          | 1.015   | M1    | oe eg 101.5% or 1 + $\frac{1.5}{100}$<br>Implied by 6090   |  |  |
|          | $6000 \times 1.015^n$ for any positive integer $n > 1$  | M1dep | oe<br>Implied by 6181.(…)  |  |  |
| 15       | 11  | A1    | If showing trials for 10 and/or 11 years,<br>must have<br>$6000 \times 1.015^{10} = 6963.()$<br>and/or<br>$6000 \times 1.015^{11} = 7067.()$ or 7068<br>If showing totals from year on year for 10<br>and/or 11 years, must have<br>(Y10) [6963.21, 6963.30]<br>and/or<br>(Y11) [7067.65, 7067.75] |  |  |
|          | Alternative method 2  |       |  |  |  |
|          | 1.015   | M1    | oe eg 101.5% or 1 + $\frac{1.5}{100}$<br>Implied by 6090   |  |  |
|          | Evaluates $1.015^{n}$ for any positive<br>integer $n > 1$<br>and<br>$7000 \div 6000$ or $1.166$ or $1.167$<br>or $1.17$ | M1dep |  |  |  |
|          | 11  | A1    | If showing trials for $n = 10$ and/or 11<br>must have<br>$1.015^{10} = [1.160, 1.161]$<br>and/or<br>$1.015^{11} = [1.177, 1.178]$  |  |  |

|    | Additional Guidance                                 |      |
|----|---|------|
|    | Values for working year on year                     |      |
|    | Y1 6000 × 1.015 = 6090                              |      |
|    | Y2 6090 × 1.015 = 6181.35                           |      |
|    | Y3 6181.35 × 1.015 = [6274.07, 6274.08]             |      |
|    | Y4 [6274.07, 6274.08] × 1.015 = [6368.18, 6368.20]  |      |
|    | Y5 [6368.18, 6368.20] × 1.015 = [6463.70, 6463.73]  |      |
| 15 | Y6 [6463.70, 6463.73] × 1.015 = [6560.65, 6560.69]  |      |
|    | Y7 [6560.65, 6560.69] × 1.015 = [6659.05, 6659.11]  |      |
|    | Y8 [6659.05, 6659.11] × 1.015 = [6758.93, 6759.00]  |      |
|    | Y9 [6758.93, 6759.00] × 1.015 = [6860.31, 6860.39]  |      |
|    | Y10 [6860.31, 6860.39] × 1.015 = [6963.21, 6963.30] |      |
|    | Y11 [6963.21, 6963.30] × 1.015 = [7067.65, 7067.75] |      |
|    | Answer 11 with no working                           | M2A1 |
|    | 1000 ÷ 90 = 11.1 Answer 11                          | Zero |

| Question | Answer   | Mark       | Comme   | nts |
|----------|--|------------|---|-----|
|          | $3y(3y^2 - 2)$ or $-3y(2 - 3y^2)$  | B2         | B1 $3(3y^3 - 2y)$ or $y(9y^2)$<br>or $-3(2y - 3y^3)$ or $-y(6)$ | ,   |
|          | Ad   | ditional G | uidance   |     |
|          | $3y(3y^2 - 2)$ or $-3y(2 - 3y^2)$ followed by incorrect further work<br>eg $3y(3y^2 - 2) = 3y^2(3y - 2)$ |            |   | B1  |
| 16(a)    | $3y(3y^2 - 2) = 3y(\sqrt{3}y + 2)(\sqrt{3}y - 2)$  |            |   | B2  |
|          | $3y(3y^2 - 2) = 9y^3 - 6y$ (checking)  |            |   | B2  |
|          | $3y \times (3y^2 - 2)$   |            |   | B2  |
|          | $3 \times (3y^3 - 2y)$   |            |   | B1  |
|          | $y3(3y^2 - 2)$   |            |   | B1  |

|       | (3x - 1)(x - 7) or $(1 - 3x)(7 - x)$   | B2         | B1 $(3x + a)(x + b)$<br>where $ab = 7$ or $a + 3b$<br>or $(a - 3x)(b - x)$<br>where $ab = 7$ or $a + 3b$ |    |
|-------|--|------------|--|----|
|       | Ade  | ditional G | uidance  |    |
|       | (3x + 1)(x + 7)  |            |  | B1 |
| 16(b) | (3x-1)(x-7)  |            |  | B1 |
| 10(0) | (3x-4)(x-6)  |            |  | B1 |
|       | (7-3x)(1-x)  |            |  | B1 |
|       | (10 - 3x)(4 - x)   |            |  | B1 |
|       | $(3x-1) \times (x-7)$  |            |  | B2 |
|       | Ignore any 'solutions' seen<br>eg $(3x - 1)(x - 7)$ in working with $\frac{1}{3}$ and 7 on answer line |            |  | B2 |

| Question | Answer   | Mark       | Commei                   | nts    |  |
|----------|--|------------|--------------------------|--------|--|
|          | Alternative method 1   |            |                          |        |  |
|          | $\sin 72 = \frac{h}{12} \text{ or } 12 \sin 72$<br>or $\cos (90 - 72) = \frac{h}{12}$<br>or $12 \cos (90 - 72)$<br>or $\frac{h}{\sin 72} = \frac{12}{\sin 90}$<br>or $11.4$              | M1         | oe<br>Any letter         |        |  |
|          | 16 × their 11.4  | M1dep      |                          |        |  |
|          | [182.4, 182.603] or 183  | A1         |                          |        |  |
|          | Alternative method 2   |            |                          |        |  |
| 17       | $h^{2} + (12 \cos 72)^{2} = 12^{2}$<br>or $h^{2} + (12 \sin (90 - 72))^{2} = 12^{2}$<br>or $\sqrt{12^{2} - (12 \cos 72)^{2}}$<br>or $\sqrt{12^{2} - (12 \sin (90 - 72))^{2}}$<br>or 11.4 | M1         | oe<br>Any letter         |        |  |
|          | 16 × their 11.4  | M1dep      |                          |        |  |
|          | [182.4, 182.603] or 183  | A1         |                          |        |  |
|          | Alternative method 3   |            |                          |        |  |
|          | 0.5 × 16 × 12 × sin 72 or 91.3   | M1         | oe eg 0.5 x 16 x 12 x si | in 108 |  |
|          | 2 × their 91.3   | M1dep      |                          |        |  |
|          | [182.4, 182.603] or 183  | A1         |                          |        |  |
|          | Ad   | ditional G | uidance                  |        |  |
|          | 2 × 16 × 12 × sin 72   |            |                          | M1M0A0 |  |
|          | $\sin = \frac{h}{12}$ or $\sin \theta = \frac{h}{12}$ (unless reco   | overed)    |                          | МО     |  |

| Question | Answer              | Mark | Comments |  |
|----------|---------------------|------|----------|--|
|          | A ∩ B'              | B1   |          |  |
| 18(a)    | Additional Guidance |      |          |  |
|          |                     |      |          |  |
|          |                     |      |          |  |

|       | (A U B)' | B1         |          |  |
|-------|----------|------------|----------|--|
| 18(b) | Ade      | ditional G | Guidance |  |
|       |          |            |          |  |

|    | Alternative method 1  |       |   |        |
|----|---|-------|---|--------|
|    | $5w \times w$ or $5w^2$<br>or 1620 ÷ 5 or 324<br>or trials a value of <i>w</i> for $5w^2$                             | M1    | oe<br>Any letter<br>eg 5 × 12 × 12 or 50 ×                        | 10     |
|    | $\sqrt{\frac{1620}{5}}$ or $\sqrt{324}$   | M1dep |   |        |
|    | 18  | A1    | A0 if –18 also given  |        |
|    | Alternative method 2  |       |   |        |
| 19 | $l \times \frac{l}{5}$ or $\frac{l^2}{5}$<br>or 1620 × 5 or 8100<br>or trials a value of <i>l</i> for $\frac{l^2}{5}$ | M1    | oe<br>Any letter<br>eg $\frac{60 \times 60}{5}$ or $80 \times 16$ |        |
|    | $\sqrt{1620 \times 5}$ or $\sqrt{8100}$ or 90   | M1dep |   |        |
|    | 18  | A1    | A0 if –18 also given  |        |
|    | Additional Guidance   |       |   |        |
|    | Answer 18   |       |   | M2A1   |
|    | 18 in working with 90 on answer line  |       |   | M2A0   |
|    | Trials for $5w^2$ or $\frac{l^2}{5}$ without answer 1   | 8     |   | M1M0A0 |

| Question | Answer  | Mark  | Comments  |  |
|----------|---|-------|---|--|
|          | Alternative method 1  |       |   |  |
|          | $h = kv^2$ or $5 = k \times 10^2$<br>or $5 \div 10^2$ or $5 : 10^2$               | M1    | oe  |  |
|          | $(k =) \frac{1}{20}$ or $(k =) 0.05$<br>or $h = \frac{1}{20}v^2$ or $h = 0.05v^2$ | A1    | oe<br>Correct value for $k$<br>or correct equation in $h$ and $v$                     |  |
|          | their $\frac{1}{20} \times 24^2$  | M1dep | oe<br>$\frac{1}{20} \times 24^2$ implies M1A1M1                                       |  |
| 20       | 28.8  | A1ft  | ft their k and M1A0M1   |  |
|          | Alternative method 2  |       |   |  |
|          | $kh = v^2$ or $k \times 5 = 10^2$<br>or $10^2 \div 5$ or $10^2 : 5$               | M1    | oe  |  |
|          | $(k =) 20 \text{ or } 20h = v^2$  | A1    | oe<br>Correct value for $k$ or correct equation<br>or correct equation in $h$ and $v$ |  |
|          | 24 <sup>2</sup> ÷ their 20  | M1dep | oe<br>24 <sup>2</sup> ÷ 20 implies M1A1M1   |  |
|          | 28.8  | A1ft  | ft their k and M1A0M1   |  |

| Question | Answer   | Mark         | Comme   | nts  |
|----------|--|--------------|---|------|
|          | Alternative method 3   |              |   |      |
|          | $\left(\frac{24}{10}\right)^2$ or $\frac{576}{100}$ or $24^2:10^2$ | M1           | oe  |      |
|          | $\frac{h}{5} = \left(\frac{24}{10}\right)^2$                       | A1           | oe<br>Correct equation in $h$                         |      |
|          | 5 × their $\left(\frac{24}{10}\right)^2$                           | M1dep        | oe<br>5 x $\left(\frac{24}{10}\right)^2$ implies M1A1 | M1   |
|          | 28.8   | A1ft         | ft their $\left(\frac{24}{10}\right)^2$ and M1A0      | M1   |
|          | Alternative method 4   |              |   |      |
|          | $\left(\frac{10}{24}\right)^2$ or $\frac{100}{576}$ or $10^2:24^2$ | M1           | oe  |      |
| 20       | $\frac{5}{h} = \left(\frac{10}{24}\right)^2$                       | A1           | oe Correct equation in $h$                            |      |
|          | $5 \div \text{their}\left(\frac{10}{24}\right)^2$                  | M1dep        | oe<br>5 ÷ $\left(\frac{10}{24}\right)^2$ implies M1A  | IM1  |
|          | 28.8   | A1ft         | ft their $\left(\frac{24}{10}\right)^2$ and M1A0      | M1   |
|          | Additional Guidance  |              |   |      |
|          | $h \alpha v^2$ with no further valid working                       |              |   | Zero |
|          | $h = kv$ or $h = kv^3$ or $h = \frac{k}{v^2}$ etc not recovered    |              |   | Zero |
|          | Up to first two marks can be awarded subsequently used             | l for correc | t working even if not                                 |      |
|          | Allow use of other letters   |              |   |      |

| Question | Answer  | Mark | Comme  | nts           |
|----------|---|------|--|---------------|
| 21(-)    | Draws $y = 3x$<br>and<br>(x =) [-0.1, 0.1] and $(x =) [1.4, 1.6]$ | B2   | B1 Draws $y = 3x$ or stat<br>$\pm \frac{1}{2}$ square tolerance for<br>Graph must be seen for<br>from 0 to 1.5 | drawing graph |
| 21(a)    | Additional Guidance   |      |  |               |
|          | Ignore any y values seen  |      |  |               |
|          | Solutions from a non-graphical method                             |      |  | B0            |
|          | Ignore other lines drawn on grid                                  |      |  |               |

| Question | Answer   | Mark | Comme   | nts  |  |
|----------|--|------|---|--|--|
| 21(b)    | Full evaluation of method and answer   | B2   | eg1 Cannot divide by $x$<br>eg2 Should have factori<br>would have also found th<br>eg3 Should have used to<br>then he would have used a<br>then he would have used a<br>then he would have complet<br>then he would have complet<br>then he would have also<br>B1 Partial evaluation<br>eg1 $x = 0$ has been omitien<br>eg2 Should have factorise<br>eg3 Should have used th<br>eg4 Should have drawn<br>eg5 Only found one solute<br>eg6 Cannot divide by ze | sed and then he<br>hat $x = 0$<br>he formula and<br>found that $x = 0$<br>a graphical method<br>found that $x = 0$<br>eted the square<br>found that $x = 0$<br>tted<br>sed<br>he formula<br>a graph<br>ttion |  |
|          | Additional Guidance  |      |   |  |  |
|          | For B2 there needs to be an evaluation that $x = 0$ has been omitted from the an |      | ethod and an indication   |  |  |
|          | x(2x + 5) = 0<br>x = 0 and $x = -2.5$  |      |   | B2   |  |
|          | Should be two solutions  |      |   | B1   |  |
|          | What about $x = 0$   |      |   | B1   |  |
|          | The answer is wrong  |      |   | B0   |  |
|          | Ignore non-contradictory further work  |      |   |  |  |

| Question | Answer   | Mark  | Comments  |
|----------|--|-------|---|
|          | Alternative method 1   |       |   |
|          | $(\frac{1}{2} \times) \pi \times 25 \times 25$<br>or $625\pi$ or $312.5\pi$<br>or [1962.5, 1964] or [981, 982]<br>or $\pi \times 12 \times 12$ or $144\pi$<br>or [452, 452.45] | M1    | oe<br>Area of circle or semicircle radius 25<br>or area of circle radius 12     |
|          | $\frac{150}{360} \text{ or } \frac{5}{12} \text{ or } 0.41(6)$<br>or 0.417 or 0.42<br>or $\frac{360}{150}$ or $\frac{12}{5}$ or 2.4  | M1    | May be seen in two steps<br>eg × 150 ÷ 360                                      |
| 22       | their $\frac{150}{360} \times \pi \times 12 \times 12$<br>or $\pi \times 12 \times 12 \div$ their $\frac{360}{150}$<br>or $60\pi$ or [188.4, 188.52]                           | M1dep | oe<br>dep on M2<br>Area of sector   |
|          | their [188.4, 188.52] (× 100)   their [981, 982] or [0.19, 0.1922]   or [19, 19.22] or [19, 19.22]   | M1dep | oe<br>dep on M3<br>their [981, 982] must be the<br>area of semicircle radius 25 |
|          | [19, 19.22]<br>and No<br>or [0.19, 0.1922]<br>and 0.2 and No   | A1    |   |

| Question | Answer   | Mark  | Comments  |
|----------|--|-------|---|
|          | Alternative method 2   |       |   |
|          | $(\frac{1}{2} \times) \pi \times 25 \times 25$<br>or $625\pi$ or $312.5\pi$<br>or [1962.5, 1964] or [981, 982]<br>or $\pi \times 12 \times 12$ or $144\pi$<br>or [452, 452.45] | M1    | oe<br>Area of circle or semicircle radius 25<br>or area of circle radius 12 |
| 22       | $\frac{150}{360} \text{ or } \frac{5}{12} \text{ or } 0.41(6)$<br>or 0.417 or 0.42<br>or $\frac{360}{150}$ or $\frac{12}{5}$ or 2.4  | M1    | May be seen in two steps<br>eg × 150 ÷ 360                                  |
|          | their $\frac{150}{360} \times \pi \times 12 \times 12$<br>or $\pi \times 12 \times 12 \div$ their $\frac{360}{150}$<br>or $60\pi$ or [188.4, 188.52]                           | M1dep | oe<br>dep on M2<br>Area of sector   |
|          | their [188.4, 188.52] × 5<br>or [942, 942.6]   | M1dep | oe<br>dep on M3   |
|          | [942, 942.6] and [981, 982] and No   | A1    | oe eg 300 $\pi$ and 312.5 $\pi$ and No                                      |

| Question | Answer   | Mark       | Comments  |
|----------|--|------------|---|
|          | Alternative method 3   |            |   |
|          | $(\frac{1}{2} \times) \pi \times 25 \times 25$<br>or $625\pi$ or $312.5\pi$<br>or [1962.5, 1964] or [981, 982]<br>or $\pi \times 12 \times 12$ or $144\pi$<br>or [452, 452.45] | M1         | oe<br>Area of circle or semicircle radius 25<br>or area of circle radius 12         |
|          | 0.2 × their [981, 982]<br>or 62.5π or [196.2, 196.4]   | M1dep      | oe<br>dep on 1st M1<br>their [981, 982] must be the<br>area of semicircle radius 25 |
| 22       | $\frac{150}{360} \text{ or } \frac{5}{12} \text{ or } 0.41(6)$<br>or 0.417 or 0.42<br>or $\frac{360}{150}$ or $\frac{12}{5}$ or 2.4  | M1         | May be seen in two steps<br>eg × 150 ÷ 360  |
|          | their $\frac{150}{360} \times \pi \times 12 \times 12$<br>or $\pi \times 12 \times 12 \div$ their $\frac{360}{150}$<br>or $60\pi$ or [188.4, 188.52]                           | M1dep      | oe<br>dep on 1st M1 and 3rd M1<br>Area of sector                                    |
|          | [188.4, 188.52] and [196.2, 196.4]<br>and No   | A1         | oe eg 60 $\pi$ and 62.5 $\pi$ and No  |
|          | Ad   | ditional G | Guidance  |
|          | Alt 3 20% of [981, 982] does not score 2nd M1 unless evaluated correctly   |            |   |

| Question | Answer   | Mark        | Comme                                   | nts             |  |
|----------|--|-------------|---|-----------------|--|
|          | Alternative method 1   |             |   |                 |  |
|          | 30 ÷ 20 or 1.5   | M1          | May be implied by correct vertical axis | ct labelling on |  |
|          | 12 ÷ 15 or 0.8   | M1          |   |                 |  |
|          | Draws block for $65 \le x < 80$<br>with height 8 small squares     | A1          | Mark intention                          |                 |  |
|          | Alternative method 2   |             |   |                 |  |
|          | 12 ÷ (30 ÷ 6) or 12 ÷ 5 or 2.4                                     | M1          |   |                 |  |
|          | their 2.4 ÷ 1.5 or 1.6   | M1dep       |   |                 |  |
|          | Draws block for $65 \le x < 80$<br>with height 8 small squares     | A1          | Mark intention                          |                 |  |
| 23(a)    | Alternative method 3   |             |   |                 |  |
|          | 12 ÷ (30 ÷ 150) or 12 ÷ 0.2 or 60                                  | M1          |   |                 |  |
|          | their 60 ÷ 7.5 or 8  | M1dep       |   |                 |  |
|          | Draws block for $65 \le x < 80$<br>with height 8 small squares     | A1          | Mark intention                          |                 |  |
|          | Alternative method 4   |             |   |                 |  |
|          | 1.5 × (30 ÷ 6) or 1.5 × 5 or 7.5                                   | M1          |   |                 |  |
|          | 12 ÷ their 7.5 or 1.6  | M1dep       |   |                 |  |
|          | Draws block for $65 \le x < 80$<br>with height 8 small squares     | A1          | Mark intention                          |                 |  |
|          | Ac   | Iditional G | uidance                                 |                 |  |
|          | Draws block for $65 \le x < 80$ with height 8 small squares 3 mark |             |   | 3 marks         |  |

| Question | Answer   | Mark | Comments                       |
|----------|--|------|--------------------------------|
|          | 10 × 4.5 or 9 × 30 ÷ 6<br>or 225 ÷ (30 ÷ 6) or 45  |      | oe<br>May be seen on histogram |
| 23(b)    | or<br>$10 \times 3.6 \text{ or } 7.2 \times (30 \div 6)$<br>or $180 \div (30 \div 6) \text{ or } 36$<br>or<br>$25 \times 2 \text{ or } 10 \times (30 \div 6)$<br>or $250 \div (30 \div 6) \text{ or } 50$<br>or<br>$34.6 \times 30 \div 6$<br>or<br>$865 \div (30 \div 6)$ | M1   |                                |
|          | 173  | A1   |                                |
|          | Additional Guidance  |      |                                |
|          |  |      |                                |

| Question | Answer  | Mark  | Comments  |  |  |
|----------|---|-------|---|--|--|
|          | Alternative method 1  |       |   |  |  |
|          | 0.5 × 8 × 9 or 36<br>or (27 – 8) × 9 or 19 × 9 or 171                     | M1    | May be seen on graph                            |  |  |
|          | $0.5 \times 8 \times 9 + (27 - 8) \times 9$ or 207                        | M1dep | M2 0.5 × (27 + 19) × 9                          |  |  |
|          | 207 and Yes   | A1    |   |  |  |
|          | Alternative method 2  |       |   |  |  |
|          | 0.5 × 8 × 9 or 36   | M1    | May be seen on graph                            |  |  |
|          | $\frac{200 - \text{their } 36}{9}$ or $\frac{164}{9}$ or 18.2             | M1dep |   |  |  |
|          | 26.2 and Yes<br>or 18.2 and 19 and Yes                                    | A1    |   |  |  |
| 24       | Alternative method 3  |       |   |  |  |
|          | 0.5 × 8 × 9 or 36   | M1    | May be seen on graph                            |  |  |
|          | $\frac{200 - \text{their } 36}{27 - 8}$ or $\frac{164}{19}$ or 8.6        | M1dep |   |  |  |
|          | 8.6 and Yes   | A1    |   |  |  |
|          | Alternative method 4  |       |   |  |  |
|          | 0.5 × 8 × 9 or 36   | M1    | May be seen on graph                            |  |  |
|          | Attempt at total distance for Beth for $26.2 \leq \text{total time} < 27$ | M1dep | eg (time 26.5s)<br>0.5 × 8 × 9 + (26.5 – 8) × 9 |  |  |
|          | Correct total distance for Beth for $26.2 \le $ total time < 27 and Yes   | A1    | eg (time 26.5s)<br>202.5 and Yes                |  |  |
|          | Additional Guidance   |       |   |  |  |
|          |   |       |   |  |  |

| Question | Answer  | Mark   | Comme                                   | nts           |  |  |
|----------|---|--|---|---------------|--|--|
|          |   |  |   |               |  |  |
|          | 342.5 or 347.5  | B1   | Allow 347.49 for 347.5                  |               |  |  |
|          | 6.35 or 6.45 or 2.55 or 2.65                                | D4   | Allow 6.449 for 6.45                    |               |  |  |
|          |   | B1   | Allow 2.649 for 2.65                    |               |  |  |
|          | their 6.35 × their 2.55 or 16.1925                          |  | Must use their lower bounds for lengths |               |  |  |
|          |   | M1   | their 6.35 must be [6.3, 6.4)           |               |  |  |
|          |   | their 2.55 must be [2  |   | 2.6)          |  |  |
| 25       | their 347.5 ÷ their 16.1925                                 | Must use their upper bound for force<br>their 347.5 bound must be (345, 350] |   | und for force |  |  |
|          |   |  |   | be (345, 350] |  |  |
|          | 21.46   | A1   | Must come from 347.5 ÷ (6.35 × 2.55)    |               |  |  |
|          |   |  | or 347.49 ÷ (6.35 × 2.55)               |               |  |  |
|          | Additional Guidance   |  |   |               |  |  |
|          | 347.49 ÷ (6.35 × 2.55) = 21.46                              |  |   | B0B1M1M1A0    |  |  |
|          | 21.4 or 21.5 does not score any marks if no working is seen |  |   |               |  |  |

| Question | Answer  | Mark | Comments  |  |
|----------|---|------|---|--|
|          | Alternative method 1 Shows that <i>CB</i> (or <i>BC</i> ) is equal and parallel to <i>DE</i> (or <i>ED</i> )  |      |   |  |
|          | $(\overrightarrow{CB} =) -(\mathbf{b} - 2\mathbf{a}) - 2\mathbf{b} - \mathbf{a}$<br>or $(\overrightarrow{BC} =) \mathbf{b} - 2\mathbf{a} + 2\mathbf{b} + \mathbf{a}$  | M1   | oe method   |  |
|          | $(\overrightarrow{CB} =) \mathbf{a} - 3\mathbf{b}$<br>or $(\overrightarrow{BC} =) 3\mathbf{b} - \mathbf{a}$   | A1   | Must see correct method for $\vec{CB}$ or $\vec{BC}$  |  |
|          | CB is equal and parallel to DE  | A1   | Must see a correct vector for first A1 and have a statement   |  |
|          |   |      | oe eg <i>CB</i> is equal and parallel to <i>ED</i>  |  |
|          | Alternative method 2 Shows that <i>BE</i> (or <i>EB</i> ) is equal and parallel to <i>CD</i> (or <i>DC</i> )  |      |   |  |
| 26       | $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>or $(\overrightarrow{CD} =) -(\mathbf{b} - 2\mathbf{a}) - (\mathbf{a} - 3\mathbf{b})$<br>or $(\overrightarrow{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>or $(\overrightarrow{DC} =) (\mathbf{a} - 3\mathbf{b}) + (\mathbf{b} - 2\mathbf{a})$ | M1   | oe method   |  |
|          | $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{CD} =) \mathbf{a} + 2\mathbf{b}$<br>or  | A1   | Must see correct method for $\overrightarrow{CD}$ or $\overrightarrow{DC}$<br>oe eg ( $\overrightarrow{BE}$ =) <b>a</b> + 2 <b>b</b> and ( $\overrightarrow{DC}$ =) - <b>a</b> - 2 <b>b</b> |  |
|          | $(\vec{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>and $(\vec{DC} =) -\mathbf{a} - 2\mathbf{b}$  |      |   |  |
|          | BE is equal and parallel to CD  | A1   | Must see two correct vectors for first A1 and have a statement  |  |
|          |   |      | oe eg <i>BE</i> is equal and parallel to <i>DC</i>  |  |

| Question | Answer   | Mark | Comments   |  |
|----------|--|------|--|--|
|          | Alternative method 3 Shows that two pairs of opposite sides are parallel   |      |  |  |
|          | $(\overrightarrow{CB} =) -(\mathbf{b} - 2\mathbf{a}) - 2\mathbf{b} - \mathbf{a}$<br>or $(\overrightarrow{BC} =) \mathbf{b} - 2\mathbf{a} + 2\mathbf{b} + \mathbf{a}$<br>or $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>or $(\overrightarrow{CD} =) -(\mathbf{b} - 2\mathbf{a}) - (\mathbf{a} - 3\mathbf{b})$<br>or $(\overrightarrow{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>or $(\overrightarrow{DC} =) (\mathbf{a} - 3\mathbf{b}) + (\mathbf{b} - 2\mathbf{a})$ | M1   | oe method  |  |
| 26       | $(\overrightarrow{CB} =) \mathbf{a} - 3\mathbf{b}$<br>or<br>$(\overrightarrow{BC} =) 3\mathbf{b} - \mathbf{a}$<br>or<br>$(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{CD} =) \mathbf{a} + 2\mathbf{b}$<br>or<br>$(\overrightarrow{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>and $(\overrightarrow{DC} =) -\mathbf{a} - 2\mathbf{b}$   | A1   | Must see correct method for $\overrightarrow{CB}$ or $\overrightarrow{BC}$<br>or $\overrightarrow{CD}$ or $\overrightarrow{DC}$<br>oe eg ( $\overrightarrow{BE}$ =) <b>a</b> + 2 <b>b</b> and ( $\overrightarrow{DC}$ =) - <b>a</b> - 2 <b>b</b>   |  |
|          | $(\vec{CB} =) \mathbf{a} - 3\mathbf{b}$<br>and $(\vec{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\vec{CD} =) \mathbf{a} + 2\mathbf{b}$<br>and $CB$ is parallel to $DE$<br>and $BE$ is parallel to $CD$  | A1   | Must see three correct vectors and have<br>two statements<br>oe eg $(\overrightarrow{BC} =) 3\mathbf{b} - \mathbf{a}$<br>and $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{DC} =) -\mathbf{a} - 2\mathbf{b}$<br>and <i>BC</i> is parallel to <i>DE</i><br>and <i>BE</i> is parallel to <i>DC</i> |  |

| Question | Answer   | Mark | Commer   | nts |
|----------|--|------|--|-----|
|          | Alternative method 4   Shows that two pairs of opposite sides are equal  |      |  |     |
|          | $(\overrightarrow{CB} =) -(\mathbf{b} - 2\mathbf{a}) - 2\mathbf{b} - \mathbf{a}$<br>or $(\overrightarrow{BC} =) \mathbf{b} - 2\mathbf{a} + 2\mathbf{b} + \mathbf{a}$<br>or $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>or $(\overrightarrow{CD} =) -(\mathbf{b} - 2\mathbf{a}) - (\mathbf{a} - 3\mathbf{b})$<br>or $(\overrightarrow{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>or $(\overrightarrow{DC} =) (\mathbf{a} - 3\mathbf{b}) + (\mathbf{b} - 2\mathbf{a})$ | M1   | oe   |     |
| 26       | $(\overrightarrow{CB} =) \mathbf{a} - 3\mathbf{b}$<br>or<br>$(\overrightarrow{BC} =) 3\mathbf{b} - \mathbf{a}$<br>or<br>$(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{CD} =) \mathbf{a} + 2\mathbf{b}$<br>or<br>$(\overrightarrow{EB} =) -\mathbf{a} - 2\mathbf{b}$<br>and $(\overrightarrow{DC} =) -\mathbf{a} - 2\mathbf{b}$   | A1   | Must see correct method<br>or $\overrightarrow{CD}$ or $\overrightarrow{DC}$<br>oe eg ( $\overrightarrow{BE}$ =) <b>a</b> + 2 <b>b</b> and   | →   |
|          | $(\overrightarrow{CB} =) \mathbf{a} - 3\mathbf{b}$<br>and $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{CD} =) \mathbf{a} + 2\mathbf{b}$<br>and $CB$ is equal to $DE$<br>and $BE$ is equal to $CD$   | A1   | Must see three correct vectors and have<br>two statements<br>oe eg $(\overrightarrow{BC} =) 3\mathbf{b} - \mathbf{a}$<br>and $(\overrightarrow{BE} =) \mathbf{a} + 2\mathbf{b}$<br>and $(\overrightarrow{DC} =) -\mathbf{a} - 2\mathbf{b}$<br>and <i>BC</i> is equal to <i>DE</i><br>and <i>BE</i> is equal to <i>DC</i> |     |
|          | Additional Guidance  |      |  |     |
|          | Choose the method that gives most marks  |      |  |     |
|          | Ignore incorrect vectors if not contradictory  |      |  |     |
|          | For parallel allow in the same direction or in the opposite direction  |      |  |     |
|          | For equal to allow = or the same as  |      |  |     |
|          | Condone incorrect notation if unambiguous<br>eg $CB = -(b - 2a) - 2b - a$  |      | M1   |     |

| Question | Answer  | Mark  | Comments   |  |
|----------|---|-------|--|--|
|          | Alternative method 1  |       |  |  |
|          | $x(x + 2)$ or $x^{2} + 2x$<br>or<br>$2x \times 4$ or $8x$<br>or<br>4(x + 2) or $4x + 8$   | M1    |  |  |
|          | $x(x + 2)$ or $x^2 + 2x$<br>and<br>$2x \times 4$ or $8x$<br>and<br>4(x + 2) or $4x + 8$   | M1dep | oe<br>eg $\frac{x(x+2) - 2x \times 4}{4(x+2)}$   |  |
| 27       | $x(x+2) - 2x \times 4 = 4(x+2)$   | M1dep | oe equation with fractions eliminated dep on M2  |  |
|          | $x^2 - 10x - 8 (= 0)$   | A1    | oe 3-term quadratic equation with terms collected  |  |
|          | $\frac{-10 \pm \sqrt{(-10)^2 - 4 \times 1 \times -8}}{2 \times 1}$<br>or $\frac{10 \pm \sqrt{100 + 32}}{2}$ or $\frac{10 \pm \sqrt{132}}{2}$<br>or $5 \pm \sqrt{5^2 + 8}$ or $5 \pm \sqrt{33}$<br>or<br>[10.744, 10.745] and [-0.745, -0.744] | M1    | oe<br>Correct for their 3-term quadratic<br>Allow correct factorisation of their 3-term<br>quadratic |  |
|          | 10.74 and $-0.74$<br>with $x^2 - 10x - 8$ (= 0) oe seen   | A1    | Must both be to 2 decimal places   |  |

| Question | Answer  | Mark                | Comments   |
|----------|---|---------------------|--|
|          | <b>Alternative method 2</b> (from $\frac{x}{4} = 1$ ·   | $+\frac{2x}{x+2}$ ) |  |
|          | $x(x + 2)$ or $x^2 + 2x$<br>or<br>(x + 2) + 2x or $3x + 2or12x + 8$   | M1                  |  |
|          | $\frac{x(x+2)}{4} \text{ or } \frac{x^2+2x}{4}$<br>and<br>$\frac{x+2+2x}{x+2} \text{ or } \frac{3x+2}{x+2}$   | M1dep               |  |
| 27       | x(x + 2) = 4(x + 2 + 2x)<br>or<br>x(x + 2) = 4(3x + 2)  | M1dep               | oe equation with fractions eliminated dep on M2  |
|          | $x^2 - 10x - 8 (= 0)$   | A1                  | oe 3-term quadratic equation with terms collected  |
|          | $\frac{-10 \pm \sqrt{(-10)^2 - 4 \times 1 \times -8}}{2 \times 1}$<br>or $\frac{10 \pm \sqrt{100 + 32}}{2}$ or $\frac{10 \pm \sqrt{132}}{2}$<br>or $5 \pm \sqrt{5^2 + 8}$ or $5 \pm \sqrt{33}$<br>or<br>[10.744, 10.745] and [-0.745, -0.744] | M1                  | oe<br>Correct for their 3-term quadratic<br>Allow correct factorisation of their 3-term<br>quadratic |
|          | 10.74 and $-0.74$<br>with $x^2 - 10x - 8$ (= 0) oe seen   | A1                  | Must both be to 2 decimal places   |

| Question | Answer  | Mark  | Commer   | nts              |
|----------|---|-------|--|------------------|
|          | Alternative method 3 (from $\frac{x}{4} - 1 = \frac{2x}{x+2}$ )   |       |  |                  |
|          | $\frac{x-4}{4}$   | M1    |  |                  |
|          | $(x-4)(x+2)$ or $x^2-4x+2x-8$<br>or $x^2-2x-8$<br>and<br>$2x \times 4$ or $8x$  | M1dep |  |                  |
|          | $(x - 4)(x + 2) = 2x \times 4$<br>or<br>$x^2 - 4x + 2x - 8 = 8x$  | M1dep | oe equation with fraction dep on M2  | ns eliminated    |
|          | $x^2 - 10x - 8 (= 0)$   | A1    | oe 3-term quadratic equa<br>collected  | ation with terms |
| 27       | $\frac{-10 \pm \sqrt{(-10)^2 - 4 \times 1 \times -8}}{2 \times 1}$<br>or $\frac{10 \pm \sqrt{100 + 32}}{2}$ or $\frac{10 \pm \sqrt{132}}{2}$<br>or $5 \pm \sqrt{5^2 + 8}$ or $5 \pm \sqrt{33}$<br>or<br>[10.744, 10.745] and [-0.745, -0.744] | M1    | oe<br>Correct for their 3-term quadratic<br>Allow correct factorisation of their 3-term<br>quadratic |                  |
|          | 10.74 and $-0.74$ with $x^2 - 10x - 8$ (= 0) oe seen  | A1    | Must both be to 2 decimal places   |                  |
|          | Additional Guidance   |       |  |                  |
|          | 10.74 and -0.74 from T & I or with no working   |       |  | 6 marks          |
|          | 10.74 or -0.74 from T & I or with no working  |       | Zero   |                  |
|          | In quadratic formula, do not allow $-10^2$ for $(-10)^2$ unless recovered   |       |  |                  |