

Surname	Centre Number	Candidate Number
Other Names		0



**GCSE – NEW**

C410UA0-1



**CHEMISTRY – Component 1:  
Concepts in Chemistry**

**HIGHER TIER**

**THURSDAY, 17 MAY 2018 – MORNING**

**2 hours 15 minutes**

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	10	
2.	10	
3.	10	
4.	8	
5.	11	
6.	5	
7.	9	
8.	9	
9.	8	
10.	11	
11.	17	
12.	12	
<b>Total</b>	<b>120</b>	

C410UA01  
01

**ADDITIONAL MATERIALS**

In addition to this paper you will need a calculator and a ruler.

**INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet.

If you run out of space, use the additional page at the back of the booklet, taking care to number the question(s) correctly.

**INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

Question 11(c) is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.

Answer **all** questions.

1. (a) The table shows some information about particles found in atoms. Complete the table.

[2]

Particle	Relative mass	Relative charge
proton	.....	+1
electron	negligible	.....
neutron	1	0

- (b) Complete the following table that shows information about atoms of some elements. [3]

Element	Mass number	Atomic number	Number of protons	Number of neutrons	Number of electrons
fluorine	19	9	9	10	.....
potassium	39	19	.....	20	19
argon	.....	18	18	22	18



- (e) The definition of an element is:

*“a substance that cannot be broken down into simpler substances by chemical methods”.*


In the 1700s a chemist named Antoine Lavoisier attempted to arrange substances in a pattern. The table shows some of the ‘substances’ which Lavoisier thought were elements. He divided the ‘substances’ into four groups. He published these groups in 1789. The modern names of some of the ‘substances’ are given in brackets.

Acid-making elements	Gas-like elements	Metallic elements	Earthy elements
sulfur	light	mercury	lime (calcium oxide)
phosphorus	caloric (heat)	copper	magnesia (magnesium oxide)
charcoal (carbon)	oxygen	nickel	barites (barium sulfate)
	azote (nitrogen)	gold	silex (silicon dioxide)
	hydrogen	iron	
		zinc	


- (i) Name **one** ‘substance’ in the table which is **not** a chemical element or compound. [1]  
 .....
- (ii) The ‘earthy elements’ are now known as compounds. Suggest why Lavoisier thought they were elements. [1]  
 .....

**BLANK PAGE**

2. (a) The following apparatus is used to show the electrolysis of water.



(i) Choose the **letter** of the graph which shows the relationship between the volume of hydrogen and the volume of oxygen formed during the process. Give the reason for your choice. [2]



Letter .....

Reason .....


(ii) Explain the **movement** of  $H^+$  ions and  $OH^-$  ions during the process. [2]

.....

.....

.....


(iii) Complete the equation by drawing diagrams to represent **all** the molecules formed. [2]




(b) The table below shows the symbols of the ions present in three electrolytes and the products formed during their electrolysis. **Complete the table.** [4]

Electrolyte	Symbol of ions present in electrolyte		Name of product formed	
	Positive ion(s)	Negative ion(s)	At the cathode (-)	At the anode (+)
molten lead(II) iodide	.....	.....	lead	iodine
aqueous copper(II) sulfate	$Cu^{2+}$ $H^+$	$SO_4^{2-}$ $OH^-$	.....	oxygen
aqueous lithium chloride	$Li^+$ $H^+$	$Cl^-$ $OH^-$	hydrogen	.....

3. (a) The diagram shows three reactions of lithium.



- (i) I Balance the symbol equation for reaction 1. [1]



- II Calculate the relative formula mass ( $M_r$ ) of lithium oxide. [1]

$$A_r(\text{Li}) = 7 \quad A_r(\text{O}) = 16$$

$$M_r = \dots\dots\dots$$


- III Describe how reaction 1 is prevented from happening when storing lithium in the laboratory. [1]

.....

.....



- (ii) I Complete and balance the symbol equation for reaction 2. [2]



- II Explain the colour seen when a few drops of universal indicator are added to the solution formed in reaction 2. [2]

.....  
.....


- (iii) Write a balanced symbol equation for reaction 3. [2]

.....


- (b) Give the chemical formula of lithium carbonate. [1]

.....

4. Iron is extracted from its ore in the blast furnace. The diagram shows the main reactions occurring in the furnace.



- (a) (i) Complete and balance the symbol equation for reaction 3. [2]



- (ii) Use this reaction to explain the term *reduction*. [1]

.....

- (b) Give the type of reaction taking place in the formation of slag. Give a reason for your answer. [2]

.....

.....

- (c) Explain how calcium oxide is formed in the furnace. [2]

.....

.....

- (d) Suggest how the cost of the process is reduced by using some of the waste gases. [1]

.....

8


5. (a) Polymers are very large molecules made when many smaller molecules join together, end to end. The smaller molecules are called monomers. The process of small monomers joining together is called polymerisation. There are two types of polymerisation.

(i) Monomer **A** undergoes addition polymerisation. Complete the table. [2]

Monomer <b>A</b>	Functional group needed for addition polymerisation	Repeating unit
$  \begin{array}{c}  \text{H} \quad \quad \text{Cl} \\  \diagdown \quad / \\  \text{C} = \text{C} \\  / \quad \quad \diagdown \\  \text{H} \quad \quad \text{H}  \end{array}  $		

(ii) I Monomers **B** and **C** can undergo a condensation reaction.

Complete the diagram by showing how these two molecules join together forming two products. [2]



II Explain, using monomers **B** and **C**, the principles of condensation polymerisation. [4]

.....

.....

.....

.....

.....

.....

- (b) When manufacturers produce soft drinks they often package the same product in different materials. Each type of disposable drink container has an environmental impact.

Scientists carried out a life cycle assessment (LCA) for three different disposable drinks' containers. The table shows some information from the life cycle assessment.

	Plastic bottle (PET)	Glass bottle	Aluminium can
Raw material(s)	petroleum	sand, sodium carbonate and limestone	bauxite
Mass of carbon dioxide emitted per container during production (g)	142	226	168
Mass of 330 ml container (g) <i>(mass impacts on truckload size and therefore fuel use)</i>	11	200	24
Recycling	25% recycled into new bottles  75% recycled into other products such as wheelie bins and eco-fleece due to degradation in properties	40% recycled into new bottles  no degradation of properties therefore can be recycled indefinitely	70% recycled into new cans  no degradation of properties therefore can be recycled indefinitely
Time to break down in the environment	400 years	400 years	80 years

Use the information from the table to state which material in your opinion has the least environmental impact.

Give **three** pieces of evidence to support your choice. [3]

.....

.....

.....

.....

.....

.....

.....

.....

C410UA01  
13

6. (a) The following tests were used to identify unknown compounds **A**, **B** and **C**.

add silver nitrate solution

add dilute hydrochloric acid,  
bubble gas given off into limewater


flame test

add sodium hydroxide solution

add sodium hydroxide solution and warm mixture,  
test gas given off with damp litmus paper

These are described below as tests **1** to **5** but **not** necessarily in that order.

The charts show the results obtained for each compound.



Deduce which test is which and hence identify compounds **A**, **B** and **C**.

[3]

**A** .....

**B** .....

**C** .....


- (b) Colourless aqueous solutions of amino acids can be separated by paper chromatography. Spots appear when the paper is sprayed with a 'locating agent'.

Examiner  
only

The table shows the  $R_f$  values for some amino acids.

Amino acid	$R_f$ value
glycine	0.25
alanine	0.40
valine	0.70
proline	0.45
serine	0.30
lysine	0.15
cysteine	0.10

A student was given the chromatogram of a mixture of two unknown amino acids.



Use the information given to identify the **two** unknown amino acids in the mixture. [2]


..... and .....

C410UA01  
15

**BLANK PAGE**



7. (a) Gareth and Caroline investigated the displacement reaction between iron filings and copper(II) sulfate solution. The equation for the reaction is as follows.



Both students carried out the following procedure.

0.56 g of iron filings were added to excess aqueous copper(II) sulfate. Once all the iron filings had reacted, the copper formed was filtered, dried and weighed accurately.

The mass of copper expected was 0.64 g.

- (i) Gareth obtained a value of 0.71 g. Suggest **one** possible reason for the higher than expected mass. State how this problem could be overcome. [2]

.....

.....

.....

- (ii) Caroline obtained a value of 0.61 g. Suggest **one** possible reason for the lower than expected mass. State how this problem can be overcome. [2]

.....


.....

.....

- (b) The students were asked to find the relative positions in the reactivity series of four unknown metals, **A**, **B**, **C** and **D**.

Gareth measured the voltage formed in a simple chemical cell. He paired metals **A**, **B** and **C** in turn with metal **D**. Metal **D** is the least reactive of the metals. The voltage formed by each pair of metals is shown in the table.

*In a chemical cell, the further apart the electrode metals are in the reactivity series the greater the voltage generated.*



Metal	Voltage (V)
<b>A</b>	0.46
<b>B</b>	1.23
<b>C</b>	2.70

Caroline carried out a series of displacement reactions. She added metals **A**, **B**, **C** and **D** to separate solutions containing the nitrate of a different metal ion.

Complete the table below to show the results that would support Gareth's evidence. [2]

Use a tick (✓) to show that a reaction occurs and a cross (✗) to show that no reaction occurs.

Metal	Metal nitrate solution			
	metal <b>A</b> nitrate	metal <b>B</b> nitrate	metal <b>C</b> nitrate	metal <b>D</b> nitrate
<b>A</b>				
<b>B</b>				
<b>C</b>				
<b>D</b>				

- (c) Suggest a reason why Gareth's is a better method than Caroline's for finding the relative positions of metals in the reactivity series. [1]

.....

.....

- (d) Metal **D** has two main isotopes,  $^{63}\text{D}$  and  $^{65}\text{D}$ .

A sample of metal **D** contains 70%  $^{63}\text{D}$  atoms and 30%  $^{65}\text{D}$  atoms.

Calculate the relative atomic mass ( $A_r$ ) of metal **D** to **three** significant figures. [2]

$A_r =$  .....

9


8. (a) (i) Calcium reacts with oxygen to form calcium oxide.

Using the electronic structures below, draw dot and cross diagrams to explain the bonding in calcium oxide. Show only outer shell electrons in your diagrams. [3]

calcium 2,8,8,2

oxygen 2,6

- (ii) Complete the diagram showing the outer shell electrons in an oxygen molecule,  $O_2$ . [2]



- (iii) Calcium oxide has an ionic structure and melts at  $2572^\circ\text{C}$ . Oxygen has a simple covalent structure and melts at  $-219^\circ\text{C}$ .

Explain the difference in the melting points of calcium oxide and oxygen. [2]


.....

.....

.....

.....

(b) The diagram shows the structure of metallic silver.



Explain why silver conducts electricity.

[2]

.....

.....

.....

9

9. Alcohols can be used as fuels. **Table 1** shows the first five members of the alcohol homologous series. The theoretical values for the energy released when alcohols are burned are also shown. The value for ethanol is missing.

Alcohol	Molecular formula	Energy released (kJ)
methanol	CH <sub>3</sub> OH	658
ethanol	C <sub>2</sub> H <sub>5</sub> OH	
propanol	C <sub>3</sub> H <sub>7</sub> OH	1894
butanol	C <sub>4</sub> H <sub>9</sub> OH	2512
pentanol	C <sub>5</sub> H <sub>11</sub> OH	3130


**Table 1**

The energies of the bonds broken and formed as alcohols burn are shown in **Table 2**.

Bond	Bond energy (kJ)
O—H	464
C—C	347
C—H	413
C—O	358
C=O	805
O=O	498

**Table 2**

The following equation shows the rearrangement of atoms as ethanol burns.




(a) Calculate the energy released for the burning of ethanol.

[5]

Energy released = ..... kJ

(b) Draw an arrow (↕) on the reaction profile to show the energy change calculated in part (a). [1]




(c) Use your answer to part (a) and the information from **Table 1** to describe the relationship between the number of carbon atoms present in an alcohol and the energy released on burning. [2]

.....

.....

.....

10. Sodium thiosulfate solution reacts with dilute hydrochloric acid forming a yellow precipitate. This reaction can be investigated using the 'disappearing cross' experiment.



50 cm<sup>3</sup> of sodium thiosulfate solution was heated in a water bath until a target temperature was reached. The flask was removed from the water bath and the actual temperature was recorded just before 10 cm<sup>3</sup> of hydrochloric acid was added. A stopwatch was started immediately. The time taken for the cross to disappear was recorded. This procedure was repeated at different temperatures. The concentrations of the acid and the sodium thiosulfate solutions were kept the same in each experiment.

The results are shown below.

Target temperature (°C)	Actual temperature recorded (°C)	Time for cross to disappear (s)	Rate $1/\text{time} \times 10^{-3}$ (s <sup>-1</sup> )
20	19	250	4
30	27	167	6
40	39	62	15
50	49	33	30
60	59	17	59


- (a) Suggest a reason for the difference between the target temperature and the actual temperature recorded for each reaction. [1]

.....

.....



- (b) Choose appropriate scales and plot the rate against the **actual** temperature recorded on the grid. Draw a suitable line. [4]



- (c) (i) The following relationship is given in many text books.

*“The rate of a reaction doubles for every  $10^{\circ}\text{C}$  rise in temperature.”*

**Use your graph** to show that this relationship is true.

[2]

- (ii) Using the relationship given in part (i) find the **time**, in seconds, it would take for the cross to disappear at 70 °C. Show your working. [3]

Examiner  
only

Time = ..... s


- (iii) At 80 °C the reaction would take less than 5 seconds. Explain why the time recorded at 80 °C would be a less accurate reading than at lower temperatures. [1]

.....  
.....

11

**BLANK PAGE**

11. (a) Most of the hydrogen used in the Haber process is obtained by reacting methane with steam.



The forward reaction is endothermic.

- (i) Explain why a high temperature and a low pressure would give the maximum yield of hydrogen. [3]

.....

.....

.....

.....

.....

.....

- (ii) Calculate the atom economy for the manufacture of hydrogen using this reaction.

Give your answer to **three** significant figures. [2]

$$A_r(\text{H}) = 1 \quad A_r(\text{C}) = 12 \quad A_r(\text{O}) = 16$$

Atom economy = ..... %

- (iii) Calculate the maximum volume of hydrogen that could be formed at room temperature and pressure from 0.16 g of methane. The volume of 1 mol of gas at room temperature and pressure is 0.024 m<sup>3</sup>.


Give your answer in m<sup>3</sup>. [3]

Volume of hydrogen = ..... m<sup>3</sup>



12. (a) The label shows the ingredients in 'Sparkling Apple Drink'.

Ingredients:  
carbonic acid, apple juice,  
sugar, glucose syrup, malic  
acid, preservative (sodium  
benzoate), artificial sweetener  
(saccharin).



A student was asked to find the concentration of carbonic acid in 'Sparkling Apple Drink'. He decided to do this by titrating the drink against sodium hydroxide solution.

- (i) He found that  $25.0\text{ cm}^3$  of 'Sparkling Apple Drink' was neutralised by  $15.0\text{ cm}^3$  of sodium hydroxide solution of concentration  $0.10\text{ mol/dm}^3$ . The relative formula mass of carbonic acid is 62.



- I Calculate the student's value for the concentration of carbonic acid in  $\text{mol/dm}^3$ .

[3]

Concentration = .....  $\text{mol/dm}^3$

- II Write this concentration as a value in  $\text{g/dm}^3$ .

[1]


Concentration = .....  $\text{g/dm}^3$

- (ii) Suggest why the concentration of carbonic acid in 'Sparkling Apple Drink' is actually less than that found by the student.

[1]

.....  
.....

(b) The flow diagram shows some reactions of ethanoic acid.



(i) Name the product with the formula  $\text{CH}_3\text{COONa}$ . [1]

.....

(ii) Write a balanced symbol equation for the reaction between ethanoic acid and magnesium. [3]

.....

(c) Dilute ethanoic acid (pH 3) reacts less vigorously with magnesium than dilute hydrochloric acid (pH 1) of equal concentration. Explain the reason for this difference in behaviour. [3]

.....

.....

.....

.....

**END OF PAPER**

**BLANK PAGE**



**BLANK PAGE**



## FORMULAE FOR SOME COMMON IONS

POSITIVE IONS		NEGATIVE IONS	
Name	Formula	Name	Formula
aluminium	$\text{Al}^{3+}$	bromide	$\text{Br}^-$
ammonium	$\text{NH}_4^+$	carbonate	$\text{CO}_3^{2-}$
barium	$\text{Ba}^{2+}$	chloride	$\text{Cl}^-$
calcium	$\text{Ca}^{2+}$	fluoride	$\text{F}^-$
copper(II)	$\text{Cu}^{2+}$	hydroxide	$\text{OH}^-$
hydrogen	$\text{H}^+$	iodide	$\text{I}^-$
iron(II)	$\text{Fe}^{2+}$	nitrate	$\text{NO}_3^-$
iron(III)	$\text{Fe}^{3+}$	oxide	$\text{O}^{2-}$
lithium	$\text{Li}^+$	sulfate	$\text{SO}_4^{2-}$
magnesium	$\text{Mg}^{2+}$		
nickel	$\text{Ni}^{2+}$		
potassium	$\text{K}^+$		
silver	$\text{Ag}^+$		
sodium	$\text{Na}^+$		
zinc	$\text{Zn}^{2+}$		

# THE PERIODIC TABLE

Group 1 2 3 4 5 6 7 0

7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4	11 <b>Na</b> Sodium 11	12 <b>C</b> Carbon 6	13 <b>Al</b> Aluminium 13	14 <b>N</b> Nitrogen 7	15 <b>O</b> Oxygen 8	16 <b>F</b> Fluorine 9	17 <b>Ne</b> Neon 10												
23 <b>Na</b> Sodium 11	24 <b>Mg</b> Magnesium 12	27 <b>Al</b> Aluminium 13	28 <b>Si</b> Silicon 14	31 <b>P</b> Phosphorus 15	32 <b>S</b> Sulfur 16	35.5 <b>Cl</b> Chlorine 17	40 <b>Ar</b> Argon 18	4 <b>He</b> Helium 2												
39 <b>K</b> Potassium 19	40 <b>Ca</b> Calcium 20	45 <b>Sc</b> Scandium 21	48 <b>Ti</b> Titanium 22	51 <b>V</b> Vanadium 23	52 <b>Cr</b> Chromium 24	55 <b>Mn</b> Manganese 25	56 <b>Fe</b> Iron 26	59 <b>Co</b> Cobalt 27	59 <b>Ni</b> Nickel 28	63.5 <b>Cu</b> Copper 29	65 <b>Zn</b> Zinc 30	70 <b>Ga</b> Gallium 31	73 <b>Ge</b> Germanium 32	75 <b>As</b> Arsenic 33	79 <b>Se</b> Selenium 34	80 <b>Br</b> Bromine 35	84 <b>Kr</b> Krypton 36			
86 <b>Rb</b> Rubidium 37	88 <b>Sr</b> Strontium 38	89 <b>Y</b> Yttrium 39	91 <b>Zr</b> Zirconium 40	93 <b>Nb</b> Niobium 41	96 <b>Mo</b> Molybdenum 42	99 <b>Tc</b> Technetium 43	101 <b>Ru</b> Ruthenium 44	103 <b>Rh</b> Rhodium 45	106 <b>Pd</b> Palladium 46	108 <b>Ag</b> Silver 47	112 <b>Cd</b> Cadmium 48	115 <b>In</b> Indium 49	119 <b>Sn</b> Tin 50	122 <b>Sb</b> Antimony 51	128 <b>Te</b> Tellurium 52	127 <b>I</b> Iodine 53	131 <b>Xe</b> Xenon 54	36		
133 <b>Cs</b> Caesium 55	137 <b>Ba</b> Barium 56	139 <b>La</b> Lanthanum 57	179 <b>Hf</b> Hafnium 72	181 <b>Ta</b> Tantalum 73	184 <b>W</b> Tungsten 74	186 <b>Re</b> Rhenium 75	190 <b>Os</b> Osmium 76	192 <b>Ir</b> Iridium 77	195 <b>Pt</b> Platinum 78	197 <b>Au</b> Gold 79	201 <b>Hg</b> Mercury 80	204 <b>Tl</b> Thallium 81	207 <b>Pb</b> Lead 82	209 <b>Bi</b> Bismuth 83	210 <b>Po</b> Polonium 84	210 <b>At</b> Astatine 85	222 <b>Rn</b> Radon 86	223 <b>Fr</b> Francium 87	226 <b>Ra</b> Radium 88	227 <b>Ac</b> Actinium 89

Key

